Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.



a SB141 F6N4

NEWSLETTER

JANUARY 1984



INTERNATIONAL WORKING GROUP

ON

FIRE BLIGHT RESEARCH



INTERNATIONAL WORKING GROUP

ON

FIRE BLIGHT RESEARCH

NEWSLETTER

from the

Plant Protection Commission

International Society for Horticultural Science

in cooperation with

U.S. Deciduous Tree Fruit Disease Workers

and

European & Mediterranean Plant Protection Organization

JANUARY 1984

United States Department of Agriculture Agricultural Research Service

Appalachian Fruit Research Station Kearneysville, West Virginia, USA



Letter from the Editor

The third international workshop was a great success and all the persons involved with its organization should be duely congratulated. During the workshop, several persons asked to be added to the membership list of the working group. So, during 1983, the membership thus increased by 32 to a total of 275. Every year, however, it is becoming increasingly more laborious to assemble, prepare, and mail our newsletter. Therefore, starting next year (1985), we need to reduce our mailing list considerably and I hereby urge everyone to read the following notice about our newsletter:

NOTICE

Working Group Membership Survey

This is your final copy of the fire blight newsletter unless the last page in this 1984 issue is completed in full and returned to me by September 1.

By the end of 1983, following 17 years after its initial occurrence in the Netherlands (1966), fire blight has conquered a considerable portion of the European Continent. For all practical purposes the disease is present west of a straight line from Toulouse, France in the south, via Strassburg and Berlin to Gdansk, Poland in the north, with a small forward bulge of about 200 km in the center, from Strassburg facing eastward.

Even though the disease was reported from Egypt in 1964, the severe outbreak in 1983 raised considerable alarm in the limited pear industry in that country. Surrounded by desserts and the Mediterranean Sea, fire blight in northern Egypt should not pose a further threat to other pome fruit producing countries.

Finally, I am delighted to announce that on October 5, 1983, the first intercontinental connection was established from Wageningen (Netherlands) to Kearneysville (USA) in regard to searching the USDA fire blight literature collection through the telephone and computer net works. Anyone who wishes to retrieve titles from this literature collection should first get in contact with Mr. Gary Lightner (304-725-3451, ext. 36) who in turn will supply you with the necessary information to make the connection.

TOM VAN DER ZWET, Secretary North American Section

Ton lander lich

International Working Group on Fire Blight Research

INTERNATIONAL WORKSHOP ON FIRE BLIGHT

HELD IN BORDEAUX, FRANCE

(SEPTEMBER 12-16, 1983)

The third International Workshop on Fire Blight was held at INRA, Pont-de-la-Maye near Bordeaux, France. Following the first two meetings in 1977 (Neth.) and 1980 (W. Germ.), this workshop was attended by 75 scientists representing 20 countries. Fifty papers or posters were presented into 6 sections dealing with different aspects of the disease: Epidemiology (2 sessions), Control, Physiology and Genetics, Breeding, Survey and action programs. Even though only 10 countries (8 in Europe) actually have fire blight, emphasis was placed on how to prevent the disease from entering into Spain, Switzerland and Italy.

Following 2 days of paper poster presentations, one day was spent making a bus trip to the area of Dax (160 km south of Bordeaux) to visit and inspect the experimental field plots partly subsidized by the AGRIMED Programme Committee (Group on Fire Blight in Pears, Apples and Related Species) of the Commission of the European Communities. Presently, this is the largest experimental plot in the warmest part of Europe where fire blight is endemic, providing suitable conditions to test pears, apples and rosaceous ornamentals for resistance to the disease, and allowing climatic and epidemiological studies in a new type of climate.

EXPERIMENTAL ORCHARD

DAK

The establishment of experimental plots in a climatic zone very different from that in northern Europe gives many opportunities for fire blight research. It offers northern European research workers the opportunity to test their selections while at the same time Mediterranean workers can test the susceptibility of their cultivars and new hybrids. It also offers researchers the opportunity to study the epidemiology and ecology of the organism under different conditions found at home. However, one of the primary objectives of this project is to determine the susceptibility of the principal fruit cultivars and ornamentals of Pomoideae in Europe.

A. HISTORY

The extension of fire blight into Europe and notably into the Aquitaine fruit-growing area in 1977-78 intiated the development of a research program for this disease. Three principal reasons justified this development: first, a total absence of data on the susceptibility of European cultivars; second, a difficulty in transposing epidemiological data developed under other conditions (agronomic, climatic); and third, an inability to adapt hybrids selected for resistence in U.S.A. to the economic and cultural conditions of Europe.

As soon as fire blight was found in the Dax area (1978), I.N.R.A. and Plant Protection Service considered the possibility of developing and maintaining experimental plots and a laboratory in that area, where two essential conditions are met: a contaminated area within a small fruit growing area and a favorable climate for the disease.

This project was initiated in 1979 with the first plantings in the experimental orchard. With the financial support of EEC it is possible to provide a service to all its members.

B. PROGRAM

- 1. To determine the susceptibility of cultivars in the collection.
- 2. To study the epidemiology of fire blight
 - evaluation of climatic risks (Billing system)
 - overwintering and dissemination of the pathogen
 - survival of bacteria on host surfaces.
- 3. To evaluate classic chemical products and new formulations. To study the efficacy and phytotoxicity (coppers) of chemicals on homogenous and susceptible hosts (Golden Delicious, Cotoneaster, Salicifolius...)
- 4. To intensify the cultivar selection process to obtain a fire blight resistant pear.

C. FINANCING

Financing is required for the rental of land and for services such as labor and materials. It is shared between the "Ministere de l'Agriculture of France (managed by Plant Protection service and the E.E.C. (managed by I.N.R.A.

D. MANAGEMENT

The planting and maintenance are done by the grower (M. DUSSARAT), using the normal farm equipment. P. LECOMTE is the experienced technician in charge of the experimental fields.

E. ENVIRONMENT

The climate is one of the hottest and wettest in France with an annual rainfall near 1 200 mm. The conditions are favorable not only for tree fruit culture but also for diseases (scab, fire blight) and weeds.

Spring is often a time of alternating wet and dry periods which make the first two years of a planting very difficult. Good water retention by the soil, however, permits tree growth to recoup any losses caused by spring drought and often assures excellent growth.

To better understand the climate of this region, it is necessary to observe the wide wariety of plant species, including numerous tropical plants which prosper here.

F. ORCHARD EXPANSION

Year	1979	1980	1981	1982	1983
На	2,6	3,5	5,5	8,3	8,5

The experimental plots are located in a 30 hectare commercial farm containing apples, pears, black currents, corn, etc...


G. EXPERIMENTAL PLAN

Today 25 experimental plots compose the plan (see accompanying plan with detail of the plots).

VERGER EXPERIMENTAL DAX 1983

Détail des plantations.

. P1	OPMEMENT : Paraconthe Cotonocate C 1			
• P2	ORNEMENT: Pyracantha, Cotoneaster, Sorbus: 1979. POIRIER: Comice, Morettini, Frinc du Mas (+ 20 ans)			900 m2
. P3	, by the table (20 alis)			500 m2
. P4	1,575,1500		9	000 m2
. P5	POIRIER : Essai variétal x Pathologie : 1981		4	600 m2
	POIRIER : Collection variétale 82 variétés : 1981	}	_	250 2
	G PORTE-GREFFE : Collection variétale : Malus, Cydonia, Pyrus : 1983	{	5	250 m2
• P6	ORNEMENT: Pyracantha, Cotoneaster: 1979-1980			655 m2
. P7	POIRIER: Hybrides: 1983		1	800 m2
• P8	POIRIER: Hybrides: 1980			800 m2
. P9	POIRIER: Hybrides: 1981	}		
. P10	POIRIER: Hybrides W1 Retour Wageningen 1980	}	7	200 m2
	Marcottière PG 1981 + hybrides poiriers 1983	}		
. P11	POMMIER: Collection de 67 variétés 1980-1981		6	700 m2
. P12	POMMIER: Hybrides: 1982		1	600 m2
. P13	POIRIER: Hybrides: 1982		4	500 m2
. P14	POIRIER : Collection surgreffée 1979-1980			3 30 m2
. P15	ORNEMENT : Collection variétale : 1980-81-82-83 Cotoneaster sal. floccosus		4	600 m2
. P 16	POIRIER-POMMIER : Collection : 1979 POMMIER : Géniteurs - Présélections : 1981 PORTE-GREFFE : collection : 1981	}		900 m2
. P17	ORNEMENT : Cotoneaster sal. floccosus Essais CEE 1981-1982-1983		6 (000 m2
. P18	POIRIER : Présélections et Géniteurs : 1981			700 m2
. P19	POIRIER: Hybrides, collection variétale: 1982-1983			000 m2
. P20	POIRIER: Hybrides W2: 1982			700 m ²
. P21	POMMIER : Golden delicious : 1982			750 m2
- P22	POIRIER : Présélections INRA : 1982			150 m2
• P23	POIRIER: Hybrides W 3: 1983			200 m2
. P24	POIRIER : Beurré Hardy : 1983			900 m2
. P25	POMMIER : Golden delicious : 1982-1983			600 m2
. Pépin	ière 1982 : Greffage des hybrides sélectionnés			40 m2
	5,		2	.70 1112



Titles of Reports and Posters Presented at the 3rd International Workshop on Fire Blight in Bordeaux, Francel

REPORTS

UNITED STATES

- Aldwinckle, H. S., J. Norelli, S.J. Schwager, and R. C. Lamb.

 Evaluation of fire blight resistance of apple cultivars and breeding new resistant cultivars.
- Beer, S. V., D. W. Bauer, and E. M. Steinberger. Studies on the mechanism of pathogenesis of Erwinia amylovora.
- Beer, S. V., J. L. Norelli, S. J. Schwager, et al. A risk assessment system for fire blight.
- Beer, S. V., J. R. Rundle, and J. L. Norelli.

 Recent progress in the development of biological control for fire blight.
- Beer, S. V., J. R. Rundle, and R. L. Wodzinski.

 Interaction between <u>Erwinia amylovora</u> and <u>E. herbicola</u>, in vitro, in immature pear fruits and in apple blossoms.
- Burr, T. J., and J. L. Norelli.

 Recent progress in chemical control of fire blight.
- Norelli, J., H. S. Aldwinckle, S. V. Beer, and R. C. Lamb.

 Differential virulence of <u>Erwinia amylovora</u> to specific apple cultivars and its implication for breeding and selection of fire blight resistant apple cultivars.
- Norelli, J. and S. V. Beer.

 Factors affecting the development of fire blight blossom infections.
- van der Zwet, T., R. L. Bell, and R. C. Blake.

 Comparative evaluation of the degree of fire blight resistance in pear cultivars and selections.
- van der Zwet, T. and P. D. Van Buskirk.

 Detection of endophytic and epiphytic <u>Erwinia amylovora</u> in various pear and apple tissues.

<u>1</u>/All reports and poster abstracts will be published in Acta Horticulturae 151 which can be ordered from: Secreatry Gerneral, ISHS, de Dreyen 6, 6703 BC Wageningen, The Netherlands.

FRANCE

- Callu, D., et al.

 Present situation of fire blight in France (1982–1983).
- Gardan, L. and Ch. Manceau.

 Persistence of streptomycin on apple and pear.
- Le Lezec, M. and J. P. Paulin.

 Shoot susceptibility of some apple cultivars in South-West France.
- Manceau, Ch., J. P. Paulin, and L. Gardan.

 Effects of streptomycin sprays on leaf microflora of pear trees.
- Paulin, J. P., R. Chartier, and P. Lecomte.

 Leaf scars inoculations of pear trees with <u>Erwinia amylovora</u>.

NETHERLANDS

- Maas Geesteranus, H. P.
 Survival of epiphytic <u>Erwinia amylovora</u> bacteria and their role in epidemiology.
- Meijneke, C. A. R.

 The new fire blight control policy in the Netherlands.
- Miller, H. J. <u>Erwinia</u> detection and its significance in survival study.

ENGLAND

- Principles and application of fire blight risk assessment systems.
- Billing, E.

 Studies on avirulent strains of <u>Erwinia amylovora</u>.
- Gwinn, D. C.

 Fire blight in perry pears and cider apples in the South-West of England.

WEST GERMANY

Mappes. D.

Trial results with a new copper formulation for the control of fire blight.

Zoller. W.

Further control experiment against fire blight in the Federal Republic of Germany.

Zoller, W.

Biochemical aspects of EPS of Erwinia amylovora.

ITALY

Calzolari. A., et al.

Surveys for the presence of fire blight in some Italian fruit growing areas.

Mazzucchi, U., C. Bazzi, G. Coti, C. Gasperini, A. Calzolari.

Detection of Erwinia amylovora contaminating apple twigs during the dormant season.

BELGIUM

Vantomme, R., et al.

Summary of the research of Erwinia amylovora.

SWITZERLAND

Grimm. R.

Actual situation of fire blight control in Zwitzerland.

SPAIN

Sampayo, M. and I. Palazon.

Prevention against fire blight in an uncontaminated country, Spain.

EAST GERMANY

Kleinhempel, H., H. Kegler, W. Ficke, and H. J. Schaeffer. Methods of testing apples for resistance to fire blight.

EGYPT

Abo El Dahab, M. R., M. A. El Goorani, H. M. El-Kasheir, et al. Severe outbreaks of fire blight in Egypt during 1982 and 1983 seasons.

POSTERS

FRANCE

- Balavoine, P., and D. Callu.

 Climate and fire blight in the North of France (1972-1983).
- Boue, H. Climate and fire blight in the Garonne Valley (1978-1983).
- Cadic, A.

 Pyracantha breeding program.
- Jacquart, C., J. P. Paulin, D. Payen, and E. Billing.
 Climatic assessment of the risks of fire blight at bloom.
- Laure, P., C. Desbons, and P. Lecomte.

 Observation on activity of pollinating insects on fire blight host plants in the DAX area.
- LeComte, P., P. Laure, and J. P Paulin.

 Climate and fire blight in the DAX area (1977-1983).
- Lespinasse, Y., and J. P. Paulin.

 Apple breeding program for fire blight resistance.
- Paulin, J. P., and G. Lachaud.

 Comparison of the efficiency of some chemicals in preventing blossom infection.
- Thibault, B., and H. P. Maas Geesteranus.

 Pear breeding for fire blight resistance.

NETHERLANDS

- Kooistra, T., and J. de Gruyter.

 Efficacy of bactericides against <u>Erwinia amylovora</u> tested on flowering Cotoneaster species.
- van der Scheer, H. A. Th.

 The experimental garden for research on fire blight at Ouwerkerk, The Netherlands.
- van Teylingen, M. and C. A. R. Meijneke.
 Applicability of the Billing system for spray warnings.

ITALY

Bazzi, C., M. Gasser, and U. Mazzucchi.
Weather analysis with Billing's spring system in relation to the potential risk of fire blight outbreaks in Italy.

Fideghelli, C., et al.

Breeding program for fire blight resistance by the Istituto
Sperimentale per la Frutticoltura.

Mazzucchi, U., C. Bazzi, G. Coti, C. Gesperini, and A. Calzolari.

Detection of <u>Erwinia amylovora</u> contaminating apple twigs during the dormant season.

BELGIUM

Deckers, T. and W. Porreye.

Chemical control of Erwinia amylovora in pear orchards.

DENMARK

Dinesen, I., E. Friis, and J. E. Olesen.

Climate and fire blight. Billing's "system I" tested under Danish conditions and computerized for operation use.

POLAND

Sobiczewski, P. Study on fire blight forecasting.

CANADA

Bonn, W. G.

Efficacy of bactericides for the control of fire blight of pears.

UNITED STATES

Bell, R. L. and T. van der Zwet.

Recent advances in the USDA pear breeding program.

PRESENT STATUS AND NEW OCCURRENCES OF FIRE BLIGHT

UNITED STATES AND CANADA

MISSOURI

We have detected streptomycin resistance in 3 orchards in West central Missouri. Eleven of twenty-eight isolates were resistant to 500 ppm streptomycin. Of these, 5 were avirulent and 6 were virulent. We do not as yet know how persistent the problem is or how extensive it is from a geographic point of view. We have the impression that the problem has had tenure of about 3 seasons.

We plan to reexamine and survey the area intensively in the 1984 spring season.

R. N. Goodman Univ. of Missouri

ILLINOIS

Fire blight was not severe in Illinois in 1983. Growers with minor amounts of the disease in limited acreages are advised to remove all infected wood beginning in mid-August. Removal at that time does not stimulate additional tree growth and the disease will not spread at this time of year. Removal of "strikes" now will reduce inoculum for 1984. Strikes are also easier to find now than during dormancy.

S. M. Ries Univ. of Illinois

NEW YORK

Fire blight was generally not severe in N.Y. State in 1983. Few of 120 pear and apple orchards surveyed had more than 50 blossom infections and only 2 had 500 or more. For the proper testing of our predictive system we need a season with more extensive and widespread infection.

S. V. Beer Cornell Univ.

UTAH

Fire blight was serious this year in Utah, especially on apples. Fire blight is usually more serious on pears, but because of weather conditions at the time of the flowering of apples we had a real epidemic. The prediction model based on 62° F mean temperature was

accurate. Within 3 days after a mean temperature of 64° F, we detected E. amylovora in apple flowers. Fire blight symptoms were first observed 10 days later. Growers who sprayed, based upon our warning, had very little disease.

S. V. Thomson Utah State Univ.

NORTH CAROLINA

In North Carolina, during the 1983 season, fire blight occurrence was very scattered, causing little or no economic damage, except were it occasionally affected young apple trees.

D. F. Ritchie N.C. State Univ.

CALIFORNIA

Fire blight in 1983 in California pear orchards occurred to the least extent in the last 12 years. This was in spite of record spring rainfall and in spite of the existence of many holdover cankers left from the previous season's severe epidemic in one district (Lake County).

Since it was predictable on the basis of low average temperature and low accumulated degree hours over 65° F that blight would be minimal in all districts in 1983, numbers of treatments applied in 1983 were also at a 12 year low. In spite of the reduced copper treatment schedule followed in 1983, fresh market quality was the worst in 12 years. Growers applying needless copper treatments during cool weather did suffer the most severe russet. However, weather alone was practically enough to knock California out of the fresh market picture in regards to russeting in 1983.

With the low incidence of blight in 1983, time spent by orchardists in the removal of old infections has been correspondingly low. Consequently the blight that did occur in 1983 still largely remains in the orchards and we will start the 1984 season with a good inoculum load, as usual.

B. Zoller
The Pear Doctor, Inc.

SOUTH CAROLINA

The 1983 season was characterized by fire blight occurring late between June 1 and June 15 on apples in South Carolina. An unusual instance of fire blight was noted on the variety 'Granny Smith' apparently in association with crown rot and shot-hole and Bostrychid beetles.

W. Miller Clemson Univ.

OREGON

The 1983 season was very mild for fire blight in the Medford pear district, presumably due to cool temperature during bloom. Two interesting incidents were observed, however. A block of 'Starkrimson' ('Red Clapp's Favorite') showed many new strikes during bloom, when temperature hadn't come near to a 60° mean. This was in the presence of many holdover cankers in the trees. In another block of 'Bosc' and 'Comice' pears, limbs without cankers were found in late spring oozing fluid through entry holes of the shothole borer, staining the bark brown. These stains were used as a guide in removing infected limbs. No holdover cankers were found, although there had been blight removed from this orchard in previous years.

- D. Sugar
- S. Oreg. Expt. Sta.

GEORGIA

No blight worth mentioning in 1983.

The following is a delayed report for 1982:

Fire blight started in early March in an apple clone we had brought up from the Old Fort Valley Station. From this start, the fire blight moved south into an adjacent block of three-year-old apple seedlings. Many of these young seedlings bloomed! Precocity was due to crabapple parentage, primarily Malus atrosanguinea. Inoculation was usually, but not exclusively, through blooms. The matter of biggest interest is that these seedlings had been challenged to $\underline{\mathsf{E}}$. $\underline{\mathsf{amylovora}}$ when very young. Possible explanations:

- 1. Failure of inoculation/reinoculation technique.
- 2. Differences between susceptibility when very young and when in 3rd leaf.
- 3. Increased susceptibility and/or screening or pathotypes in open blooms.

The inoculum spread through apples scattered over 20 acres wherever susceptible genotypes existed and ruined several promising selections that had not been tested before. It got all over 4- and 5-year-old trees of 'Mollie's Delicious' as usual. I keep hoping to find a resistant sport in 'Mollie's Delicious' but then that hope hasn't worked out in 'Bartlett' too well, has it?

J. M. Thompson
S.E. Fruit & Tree Nut
Research Station

ONTARIO

This year less fire blight was observed than 1982, also a light disease year. In contrast to previous years, the disease was mostly restricted to pears. Weather conditions were nt exactly favorable for blight in 1983. An unusual occurrence of fire blight was observed in the Niagara region of Ontario this past summer. In mid-July, pear fruit were found to be oozing with bacteria; however, there was little shoot or sucker blight and old cankers did not appear to be active. This fruit blight phase continued trough August and into September. The most susceptible cultivars were 'Flemish Beauty', 'Clapp Favorite', and 'French Bartlett'. Weather conditions leading up to the first observations in July were a cool and wet May, a normal June, and a hot and dry July.

W. G. Bonn Agr. Can. Res. Sta.

NOVA SCOTIA

A severe outbreak of fire blight of apples occurred on several cultivars of one-year-old whips in a local nursery. This is the first authenticated identification (confirmed by Harrow Research Station) of fire blight on apples in Nova Scotia. Until now, fire blight on apples had been confirmed to a few reports of the presence of typical fire blight cankers. The canker phase often occurs on pears and one outbreak was observed on five trees in 1983.

R. G. Ross Agr. Can. Res. Sta.

OTHER COUNTRIES

MEXICO

Fire blight has been reported from the following hosts in these regions:

Regions	Host
State of Morelos	Pear and Crataegus
State of Mexico	Pear, Apple and Quince
State of Zacatecas	Pear and Apple
State of Chihuahua	Pear and Apple
State of Puebla	Pear and Apple
State Of Midroacan	Pear

L. Fucikovsky Centro de Fitopath. Chapingo

NETHERLANDS

Spring 1983 has been very unfavourable for the development of the disease in The Netherlands. It was cold and wet until after the blossoming period of pears, apples and hawthorns, resulting in very few blossom-infections in these genera. The temperature rose, however, in the second half of June, causing a lot of blossom infection in the wide-spread broad-leaved Cotoneaster shrubs, which were just blossoming at that time. Later, shoot infection in other host plant genera resulted from this mass inoculum.

Tracing and roguing of infected plants in 20 protected area's as well as in a number of special cases (e.g. in zones around orchards) outside these regions went on continuously. In some of these protected areas and zones scarcely any hawthorn or <u>Cotoneaster</u> is left after several years of this type of action. Remaining hawthorns have to be cut back regularly in order to prevent blossoming. Planting of the most susceptible hostplants (<u>Crataegus</u>, <u>Stranvaesia</u> and the broad-leaved <u>Cotoneaster</u> species) will be forbidden in these area's starting in 1984.

C. A. R. Meijneke Plant Protect. Serv.

WEST GERMANY

In the northern part of Germany, the weather was extremely dry and hot during the summer time. Infections were very low; only in nurseries with artificial irrigation, severe attack mainly on ornamentals (especially Cotoneaster species) could be observed. With the exception of Bavaria in the south, fire blight has been spreading considerably, in particular Cotoneaster salicifolius floccosus was heavily affected.

W. Zoller Biol. Bundesanstalt Heikendorf In 1983, incidence of fire blight was again very low in southwestern Germany. However, the disease is still spreading.

E. Seemuller Biol. Bundesanstalt Dossenheim

ENGLAND

Hawthorns were again affected in some areas largely because of rain dispersal of inoculum during bloom plus progression of 1982 infections. Where pears were affected, this was usually related to progression of undetected 1982 infections or proximity to infected hawthorns; some summer blossom infections were seen. Apple blossom and shoot infections were largely confined to late flowering cider apple cultivars; weather was cool and wet during most of the apple blossom period.

E. Billing
East Malling Res. Sta.

FRANCE

1. General tendencies

In spring, weather has been generally cool and wet, with short bloom period, especially for pears and apples. Summer (particularly July and August) has been more favourable to fire blight, with extremely warm and stormy periods in several places.

2. Fire blight in previously contaminated area

- noticeable extension in the North focus which extends in Haute Normandie (South of the previously known zone) on cider apple (for the first time in France).
- the generalization along the Rhine Valley on Cotoneaster (mainly).
- slight increase of the contaminated area in the Garonne Valley, both East- (Agen) and Westward (Langon). Isolated cases in spots in the Dordogne and Lot Valleys.

3. New zones

- a) Two important new areas with fire blight on pears ('Passe-Crassane') have been detected in summer: Orleans, where several ha of 'Passe Crassane' have been uprooted after fire blight has been recorded (on secondary blossoms in June). In Ile de France (20 km North of Paris) where ll has of orchard ('Passe-Crassane') were found very severely affected in August. The stormy weather (June-July) very warm, is probably responsible for a rapid extension with these zones. and
- b) Several records in autumn on ornamentals in new places:
 - Maine et Loire on Pyracantha, on a single hedge
 - Charentes (South of Angouleme) in nursery plant material
 - Limoges on ornamentals.

J. P. Paulin INRA, S.P.V.

NEW ZEALAND

No new occurrences of fire blight. Minor isolated outbreaks of twig blight on apples and <u>Cotoneaster</u>. More prevalent in Hawkes Bay then in Nebon and Central Otago orchards.

C. N. Hale D.S.I.R., P.D.D.

SWITZERLAND

So far no fire blight has been detected in Switzerland. Spring 1983 was wet and cold and the summer hot and dry. The rigorous quarantine measures were continued, also the efficient inspection service for export nurseries.

In 1983, about 250 samples of suspected plants were tested for fire blight in the laboratory of bacteriology. If bacterial pathogens were found, mostly Pseudomonas syringae could be identified.

R. Grimm Swiss Fed. Res. Sta.

ITALY

Up until now, no cases of fire blight have been found in Italy.

C. Bazzi Instit. Patol. Veg.

ROMANIA

Fire blight has not been found in Romania.

V. Severin Res. Inst. for Pl. Prot.

GREECE

Extensive surveys were conducted in nurseries and orchards in fruit tree growing areas. No fire blight symptoms were observed.

P. G. Psallidas Ben. Phytopath. Inst.

SPAIN

Until now, no cases of fire blight have been found in Spain.

M. C. Noval Alonso I.N.I.A., Prot. Veg.

PORTUGAL

Fire blight has not yet been found in Portugal.

J. M. S. Martins Est. Agron. Nac.

IRELAND

Fire blight has not been recorded in Ireland.

P. F. Walsh Dept. of Agric.

SWEDEN

No fire blight has been found. A survey is carried out every summer in the southern parts of Sweden. Import restrictions; no fire blight hosts are allowed; inspections in the nurseries.

M. Graberg Nat. Board of Agric.

NORWAY

Fire blight has still not been observed in Norway. If the disease should be brought into the country, for example with imported nursery products, one fears that the climatic conditions will not be an obstacle to its establishment in certain areas. Consequently, importation of the chief host plants of fire blight, from countries considered to be contaminated by the disease, is prohibited.

H. Roed Norw. Pl. Prot. Inst.

AUSTRALIA

Fire blight has not been found within Australia and stringent plant quarantine measures apply to prevent its entry.

D. N. CartwrightS. Austr. Dept. of Agric.

DETAILS ON CURRENT FIRE BLIGHT RESEARCH

REPORTED FROM SOME UNIVERSITIES AND EXPERIMENT STATIONS

NETHERLANDS .

The efficacy of six compounds in one spray after inoculation with \underline{E} . amylovora (10^7 cells/ml), were tested on flowering Cotoneaster dammeri 'Coral Beauty'. In the preventive trials, Kasumin 25% wp (kasugamycin). MBR 10995 (50% experimental bactericide), Plantomycin (streptomycin), Tri-Miltox forte NC (Mancozeb, coppersulphate, copperoxychloride, coppercharbonade) and Koper Bayer (copperoxychloride) was determined exclusively by the copper content of the treatment and not by the compound. In the curative test, Kasumin has shown to be more active than the other investigated products namely Plantomycin.

In trials under natural infection conditions on <u>Cotoneaster salicifolius</u> cv. <u>floccosus</u> repeated treatments with kasumin, <u>Tri-Miltox</u> forte NC and Kiper Bayer gave sufficient control of flower infection. In the advised dosage, Copac E was not effective.

In phytotoxicity trials Copac E and Koper Bayer, applied a the same copper dosage, were equally phytotoxic on Malus 'Golden Hornet', Stranvaesia davidiana, Cotoneaster bullatus and seedlings of Malus communis and Sorbus aucuparia. In the advised dosage, Tri-Moltox forte NC and Kasumin 25% wp were less phytotoxic than Koper Bayer on Malus 'Golden Hornet' and Cotoneaster bullatus. On pear, 'Conference' and 'Clapp's Favorite', Kasumin 25% wp gave too much leaf damage.

T. Kooistra Wageningen

Trials on the survival of <u>E</u>. <u>amylovora</u> bacteria on plant part surfaces and on dead materials were continued to investigate the necessity of desinfection. Except for the ooze itself, survival of free living bacteria or bacteria clustered in strands does not extend a period of 24 hours.

H. P. Maas Geesteranus Wageningen

WEST GERMANY

Because of toxicological aspects, the further work with the very effective compound CGA 78 039 had to be stopped. Alternatively, spray experiments were undertaken with Copac E, a product with a very low Cu concentration. First results showed a good effect against artificial shoot infections on pear (up to 70% efficiency) but low effect against blossom infections on Cotoneaster salicifolius floccosus. Breeding for resistance in the highly susceptible Cotoneaster species C. salicifolius and C. watereri were continued. Five from 15 lines showed a low degree

of susceptibility and will be further worked with. The resistance studies with an assortment of pome fruit varieties will be continued on the test plot near Bornhoved.

The weather system of BILLING was tested in the north of Germany in the fruit free area of the "alte Land" and will be continued in the next year (Baumm, Univ. of Hamburg). A comparison of different epidemiological methods has been undertaken in the south of Germany in Baden-Wurttemberg (Schili, University of Stuttgart-Hohenheim)

W. Zoller Heikendorf

ENGLAND

From studies reported on other Enterobacteriaceae it appears that some non-capsulated phage-resistant variants of \underline{E} . $\underline{amylovora}$ lack the enzyme UDP-Gal-4-epimerase.

Billing's risk assessment systems gave useful warning of risks in 1982 and 1983 and reflected subsequent events well.

E. Billing
East Malling

Cell surface contact was required between virulent bacteria and slices of immature pears to induce leakage of pear electrolytes and maintain bacterial growth. Enhanced production of outer membrane component of bacteria reduced loss of pear electrolytes, and inhibited disease in inoculated susceptible apple plants.

R. C. Hignett East Malling

BELGIUM

Further investigations are made into the role of the honeybee in the dissemination of fire blight. Besides research on the spread of the bacteria by the honeybee while visiting flowers, the overwintering of the bacteria in a beehive is investigated.

O. Van Laere Res. Sta. for Nemat.

FRANCE

Studies on genetical aspects of pathogenicity are presently beginning in Institut National Agronomique (Pathologie Vegetale - Prof. Coleno) by A. Kotoujansky and J. Vanneste, in cooperation with INRA-Angers.

J. P. Paulin Angers

ITALY

Testing of plant material imported from abroad (Ministerial Decree, 23/XII/1983) for presence of E. amylovora.

Breeding program for fire blight resistance by the Istituto Sperimentale per la Frutticoltura (Rome, Italy).

C. Bazzi Bologna

SPAIN

Regular screenings of fruit-growing areas and forest with abundant hawthorns.

Systematic samplings of imported plant material likely to carry Erwinia amylovora.

Conducting various kinds of studies: fire blight risk evaluation in Spanish fruit-growing areas, bactericide phytotoxicity studies on vegetation and along the fertilization process, studies on the sensitivity of native varieties to the disease.

M. C. Noval Alonso Madrid

NEW ZEALAND

Survival of epiphytic <u>Erwinia amylovora</u> on natural apple and pear fruits and methods for disinfestation. This project is partly funded by the New Zealand Apple and Pear Marketing Board in conjunction with the Department of Scientific and Industrial Research. The aim of the project is to provide export fruit which does not carry <u>Erwinia amylovora</u>.

C. N. Hale Auckland

NEW YORK

Mutants of Erwinia amylovora that are altered in pathogenicity towards rosaceous hosts, express differential virulence towards apple cultivars or fail to induce the hypersensitive reaction in tobacco, are being produced by transposon mutagenesis. Preliminary results indicate that mutants in the three functions have been produced by use of Tn5 in the vehicle pJB4JI. DNA from a mutant deficient in pathogenicity has been isolated and a restriction endonuclease fragment containing the transposon has been identified preparatory to its cloning for use as a probe for isolation of the homologous wild-type gene. A transformation procedure for E. amylovora has been developed by optimizing buffers, pH, additional reagents, temperatures and times of treatment. Its efficiency is considered more than sufficient for planned complementation studies.

Data were collected which indicate that biological control methods for fire blight are possible. The most encouraging technique involves the application of log-phase suspensions of Erwinia herbicola, a nonpathogenic endophytic and epiphytic bacterium. Laboratory and field methods were developed for identifying those strains of E. herbicola that are particularly effective in suppressing the development of fire blight of apple blossoms. Of several hundred strains tested by various means, several consistently provided control of blossom infection of apple that is statistically equivalent to that provided by streptomycin. In addition to strain, the concentration, time of application and the physiological state of applied bacteria were found to affect efficacy.

Initially, bacteriocinogenicity was considered in selecting strains of <u>E. herbicola</u> strains for biological control studies. However, field and laboratory experiments indicated that other mechanism are responsible for control of fire blight by <u>E. herbicola</u>. A mechanism based on the utilization of endogenous organic nitrogenous compounds by <u>E. herbicola</u> is supported by the strongest evidence. One bacteriocin (Herbicolacin 112Y) was found to specifically inhibit all wild-type strains of <u>E. amylovora</u> tested. Its mode of action was studied and it was partially purified and identified as a biologically unique compound composed of several amino acid derivatives. Genes for its production (or regulation) were determined to reside on a large indigenous plasmid, based on analysis of DNA from transposon-induced non-bacteriocinogenic mutants, and the transfer of bacteriocin producing ability by conjugal transfer of the wild-type plasmid.

Work continued on the development of a system for the evaluation of the relative risk of the occurrence of fire blight in particular pear and apple orchards in a given growing season. The system takes into account factors peculiar to individual orchard, which affect host susceptibility and the availability of inoculum, and weather conditions that affect the dissemination and multiplication of the fire bight pathogen Erwinia amylovora. A working model was developed that relates weather risk and orchard risk to determine overall risk. Data were collected from 100 orchards in 1982 and 120 orchards in 1983. These data are being used to test and possibly revise the model. Plans have been made to collect additional data in 1984.

Data relating environmental conditions to the development of fire blight blossom infection under precisely controlled conditions indicated that temperature, relative humidity and inoculum dose significantly affect the rate of development of infection.

S. V. Beer Ithaca

UTAH

<u>Erwinia amylovora</u> was detected almost exclusively on the pistils of epiphytically colonized pear, apple and hawthorn flowers.

S. V. Thomson Logan

ONTARIO

Current fire blight projects include those on epidemiology involving weather monitoring and disease control using experimental compounds.

G. W. Bonn Harrow

MEXICO

Epidemiological studies on fire blight in the State of Morelos (Blooming of pear in Sept. Oct. Nov. Dec. Jan. and Feb.). Infection starts in Jan. and Feb.

MISCELLANEOUS NEWS

During late May 1983, Dr. van der Zwet (WVa) was invited to spend 10 days in Egypt, upon invitation by the Ministry of Agriculture, to diagnose and confirm the presence of fire blight on pears in the lower Nile Delta. Upon the first introduction of fire blight (origin unknown) in Egypt in the early 1960s, the disease apparently increased slowly and became widespread and yet unnoticed until recent years. In 1983, unusually late rainstorms during and after bloom (April) appeared to be the main factor initiating primary infection throughout the lower part of the Delta region. Fire blight was found widespread and quite severe just south of Alexandria in Behera Governarate, the principal region of low chilling pears, predominantly of the 'Le Conte' variety. The prevailing dessert climate between April and November is expected to be very effective in reducing further spread of the disease.

Earlier in May, Dr. Zeller visited the University of Alexandria, Egypt. On several trips in the pear orchards he observed heavy fire blight infections, mostly on blossoms on the variety 'Le Conte'. Sometimes a strong ooze production was noted. Accordingly, an Egyption student shall prepare her Masters thesis on fire blight in Heikendorf at the Biologische Bundesanstalt.

During the summer of 1983, Dr. Bonn (Harrow), spent 3 months in the laboratory of Dr. Paulin at INRA, Angers in France. His interests were serology and fire blight forecasting, two areas of research where the INRA laboratory has expertise. He enjoyed the hospitality of the French very much.

In September 1983, Dr. Sherm Thomson, Utah State University, Logan, visited Mt. Albert Research Station in New Zealand and presented a seminar on fire blight in the USA.

In September 1983, following the workshop on fire blight in Bordeaux, Dr. van der Zwet visited fruit research stations in Nyon and Wadenswil, Switzerland, in Lana and Laimburg, Italy (South Tirol), and in Bavendorf, Dossenheim (BBA), and Limburgerhof (BASF), West Germany. En route, he presented two invitational seminars in German: "Feuerbrand und die Moglichkeiten seiner Bekampfung" at the Federal Research Station for Fruit Growing in Wadenswil and "Stand und Massnahmen der Feuerbrand Wirtschaft und fur Resistenter Apfel und Birnensorten im Nord Amerika" at the Institute for Plant Protection of the University of Hohenheim near Stuttgart.

Presently, Dr. Klement (Budapest) is an invited visiting professor in the Institute fur Pflanzenpathologie und Pflanzenschutz der George August Universitat in Gottingen, West Germany (Sept. 1983-Sept. 1984).

Currently, Dr. Steve Beer (Cornell) is on a 3-month sabbatic at the Microbiologisches Institut, Swiss Federal Technical University (ETH), Zurich. He is working on molecular biology of bacteria and lecturing on phytobacteriology. Some cooperative work on fire blight with Dr. Richard Grimm in Wadenswil (Jan. - Sept. 1984).

Mrs. Jacob Conceicao, of the Portuguese Plant Protection Service, is now specializing in Plant Bacteriology. She will be responsible for the survey of bacterial plant diseases in her institution, thus becoming one more person in this country whose professional activity is concerned with the fire blight problem.

Joel Vanneste, presently training in genetics at I.N.A. Paris, will be included in the Research Staff of Angers Pathology Station (INRA) after he completes his thesis (2 years).

Dr. Douglas W. Dye, Head of the Bacteriology Section, Plant Disease Division, D. S. I. R., retired on December 16, 1983. Work on fire blight is now under the direction of Dr. Chris Hale, who has succeeded Dr. Dye as head of the Bacteriology Section.

Dr. Eve Billing retired from her post at East Malling Research Station on 31 May 1984 on reaching the age of 60. This is now the rule. At present she is a part-time consultant to the Ministry of Agriculture Fisheries and Food advising on the application of risk assessment systems. Her present aim is to round off work on the systems and on host parasite relationships and write it up. Limited writing and lecturing on other aspects of plant bacteriology is also proposed.

At present there is no sign of a replacement full-time worker on fire blight at EMRS but hopefully some work will continue there and elsewhere in spite of current financial restraints in agricultural research.

FUTURE MEETINGS

April 4

The Association of Applied Biologists in the U.K. is organizing a meeting on fire blight in London. It is primarily concerned with experience of the disease in England. No one is now working full-time on fire bight but about 30 people in research institutes, universities and the Agricultural Development and Advisory Service have spent a small proportion of their time on some aspect of the disease in recent years.

July 16-18

National Congress of the Mexican Phytopathological Society in San Luis Potosi. Mexico.

August 12-16

Annual meeting of the American Phytopathological Society in conjunction with the Canadian Phytopathological Society; University of Guelph, Guelph (Ontario), Canada.

June 2-7, 1985

Sixth International Conference on Plant Pathogenic Bacteria; Center of Adult Education, University of Maryland, College Park, MD. Contact: Dr. E. L. Civerolo, USDA Fruit Laboratory, Room 111 Bldg 004, BARC-W, Beltsville, MD 20705.

August 11-15, 1985

Annual meeting of American Phytopathological Society; MGM Grand Hotel, Reno, Nevada.

August 11-20, 1986

22nd International Horticultural Congress, University of California, Davis.

September 1986

4th International Workshop on Fire Blight, to be held at Cornell University (Ithaca) and at the N.Y. State Agricultural Experiment Station (Geneva) with tours arranged afterwards to surrounding experiment Stations (Harrow, Beltsville, Kearneysville). First announcements will be mailed in September 1984.

NEW THESES AND DISSERTATIONS ON FIRE BIGHT

Alberto Mendoza Herrera

"Identificacion y Evaluacion de Problemas Fitopatologicos del Peral (Pyrus communis L.) en el Elido Ocoxaltepec", Ocuituco, Morelos. M.S. Thesis, Colegio De Postgraduados, Chapingo, Mexico, 1983.

Pascale Garret

"Contribution a la mise au point d'un test de sensibilite a <u>Erwinia amylovora applicable a la selection in vitro de plants resistantes au Feu Bacterien", 43 p, Ecole Nationale Superieure Agronomique Femine, Rennes, Septembre 1983.</u>

Joel Vanneste

"Etude du plasmide d'<u>Erwinia amylovora</u>-Son roe dans le pouvoir pathogene." Diplome d'Etude Approfondie, Fac. Sci. Univ. Paris XI, 1983.

Giuseppe Coti

"Messa a punto di un metodo per il rilevamento dei germi di <u>Erwinia amylovora</u> su materiale asintomatico." M.S. Thesis, Univ. of Bologna, Italy, 1984.

Alia Shoeib

"Studies on fire blight disease of pears." M.S. Thesis, University of Alexandria, 1984.

Rizaldo Bayot

"Role of Flagella Motility in Apple Blossom Invasion and Tactic Response to Various Plant Nectar Extracts by Erwinia amylovora." Ph.D. Dissert., Univ. Illinois, 1984.

Michael Klopmeyer.

"Motility and Chemotaxis of <u>Erwinia herbicola.</u>" Ph.D. Dissert., Univ. of Illinois, 1984.

FIRE BLIGHT LITERATURE RECEIVED DURING 1983

(Not listed in USDA Agriculture Handbook 510, the Additional Bibliography or Previous Newsletters)

UNITED STATES

UN]	TED STATES	
	I-A-53	Aldwinckle, H. S. and S. V. Beer. 1982 Apple. pp. 642-645. <u>In</u> McGraw-Hill Encyclopedia of Science and Technology, McGraw Hill, New York.
	I-A-54	van der Zwet, T. and S. V. Beer. 1982 Fire blight, the scourge of the pear industry. pp. 375-392. In The Pear, Hortic Public., Gainesville, Fla.
	I-A-55	Aldwinckle, H. S. and S. V. Beer. 1979. Fire blight and its control. Hortic. Rev. 1:423-474.
	111-214	Giddix, L. R., F. L. Lukezic, and E. J. Pell. 1981. Effect of light on bacteria [Erwinia amylovora] -induced hypersensitivity in soybean. Phytopath. 71:111-115.
	111-215	Leong, S. A., and J. B. Neilands. 1982. Siderophore production by phytopathogenic microbial species. Arch. Biochem. Biophys. 218:351-259.
	III-216	Gantotti, B. V. and S. V. Beer. 1982. Plasmid borne determinants of pigmentation and thiamine prototrophy in Erwinia herbicola. Jour. Bact. 151:1627-1629.
	III-217	Steinberger, E. M. and S. V. Beer. 1983. Transposon mutagenesis of Erwinia amylovora with transposon TN-5. Phytopathology 73:1347.
	III-218	Bauer, D. W. and S. V. Beer. 1983. Transformation of <u>Erwinia amylovora</u> with the plasmid PBR-322. Phytopathology 73:1342.
	III-219	Sijam, K., A. L. Karr, and R. N. Goodman. 1983. Comparison of the extracellular poly saccharides produced by <u>Erwinia amylovora</u> in apple tissue and culture medium. Physiol. Plant Pathol. 22:221-232.
	IV-97	Bauer, D. W. and S. V. Beer. 1983. Evidence that the activity of a putative necro toxin of Erwinia amylovora is due to the action of inorganic salts. Phytopathology 73:362.
	IV-98	Beer, S. V., T. M. Sjulin, and H. S. Aldwinckle. 1983. Amylovorin induced shoot wilting lack of correlation with susceptiblility to Francisco amylovora

susceptiblility to

Phytopathology 73:1328-1333.

Erwinia

amylovora.

with

- IV-99

 Beer, S. V. and J. R. Rundle. 1983.

 Suppression of <u>Erwinia amylovora</u> by <u>Erwinia herbicola</u>
 in immature pear fruits. Phytopath. 73:1346.
- Van Alfen, N. K. 1982.

 Wilts: Concepts and mechanisms [Erwinia amylovora
 Corynebacterium michiganense]. p.459-474 In,
 Phytopathogenic Prokaryoles, Academic Press, New York.
- IV-101 Ferree, D. C., M. A. Ellis, and F. R. Hall. 1983.

 Tree loss due to fire blight infection of rootstocks and interstems in Ohio apple orchards. Compact Fruit Tree 16:116-120.
- IV-102 van der Zwet, T. 1983.

 Occurrence of fire blight in commercial pear seedling rootstocks following budding with symptomless scionwood. Phytopathology 73:969.
- V-42 Purcell, A. H. 1982.
 Insect vector relationships with prokaryotic plant pathogens. Ann. Rev. Phytopath. 20:467.
- VII-50

 Beckman, J. S. and J. N. Diedow. 1982.

 Studies on the bactericidal action of hydro quinone oxidation catalyzed by peroxidase. Plant Physiol. 69:22.
- VII-51 Goodman, R. N. 1981.

 Ultrastructural alterations of xylem vessel walls by pathogenic bacteria. Agronomie 1:822.
- VII-52 Goodman, R. N. 1983. Fireblight - A Case Study. Biochem. Plant Path. 4:45-63.
- VIII-23 Lombard, P. B., P. H. Westigard, J. G. Strang,
 R. Allen, and D. N. Joy. 1982.

 Effect of nitrogen and daminozide on shoot growth for pear Psylla suppression and on 'Barlett' pear performance. HortScience 17:668-669.
- IX-254

 Keil, H. L. 1976.

 Chemical control of fire blight [Erwinia amylovora] in pear by means of tree trunk injection. Proc. 3rd Workshop on Fire Blight Res. pp.97-99.
- IX-255 Goodman, R. N. 1981.

 Understanding fire blight and its control with streptomycin. Mich. State Hort. Soc. Ann. Rep. 111:50-52.
- IX-256 Covey, R. P. 1982.
 What about fire blight? Proc. Wash. State Hort.
 Assoc. Ann. Mtg. 78:171-173.
- IX-257 van der Zwet, T. 1983.

 When can we expect predictable control of fire blight? Compact Fruit Tree 16:127-138.

Norelli, J. L. and T. J. Burr. 1984. IX-258 Chemical control of fire blight of apple during bloom. Amer. Phytopath. Soc., Fung. & Nemat. Tests 39:16-17. Norelli, J. L. and T. J. Burr. 1984. IX-259 Evaluation of copper materials as early season spray to reduce primary inoculum of Erwinia amylovora in the orchard. Amer. Phytopath. Soc., Fung. & Nemat. Tests 39:17. Waterworth, H. and C. N. Clayton. 1982. XI-282 Fire blight in Bradford pear. HortScience 17:789. Aldwinckle, H. S., J. L. Preczewski, and S. V. Beer. XI-283 1978. Differences in response of apple (Malus pumila) cultivars to inoculation with Erwinia amylovora. Proc. 4th Int. Conf. Plant Path. Bacteria pp. 505-512. XI-284 Westwood, M. N., P. B. Lombard, and H. O. Bjorstand. 1976. Performance of 'Bartlett' pear on standard and old home X Farmingdale clonal rootstocks. Amer. Soc. Hort. Sci. 101:161-164. Cummins, J. N. 1983. XI-285 new dwarfing, apple rootstock from West Germany. Compact Fruit Tree 16:98 XI-286 Westwood, M. N. and P. B. Lombard. 1983. Pear rootstocks: present and future. Fruit Var. Jour. 37:24-28. XI-287 van der Zwet, T. 1983. Effect of plant height and Erwinia amylovora inoculum concentration on fire blight infection in pearch seedlings. Phytopathology 73:969-970. XI-288 Lane, R. P. and J. W. Daniel. 1983. Cultivar 'Spalding', a new home orchard pear. HortScience 18:166. XI-289 Sherman, W. B., C. P. Andrews, P. M. Lyrene, and R. H. Sharpe. 1982. 'Flordahome'-A pear for homeowners in north and central Florida. Circular, Agric. Exper. St., Fla. Univ. No. S-287, p.4. XI-290 Way, R. D., K. G. Livermore, and H. S. Aldwinckle. 1982. 'Early Cortland' apple. HortScience 17:990.

Lamb, R. C., H. S. Aldwinckle, and D. E. Terry. 1983.

Food and Life Sci. Bull. 102, 2pp.

'Freedom'-A new disease-resistant apple. New York's

XI-291

XI-292 Cummins, J. N., H. S. Aldwinckle, and R. E. Byers. 1983. 'Novole' Apple. HortScience 18:772-774. XI-293 Way, R. D., K. G. Livermore, and H. S. Aldwinckle. 1982. "Early Cortland' and 'Geneva Early' apples. York's Food and Life Sci. Bull. 99, 3pp. XI-294 Norelli, J. L., H. S. Aldwinckle, and S. V. Beer. 1984. Differential host-pathogen interactions among cultivars of apple and strains of Erwinia amylovora. Phytopathology 74:136-139. EL-80 Thomas, C. S. and J. H. Hart. 1983. Fire blight of trees and shrubs [caused by the Bacterium Erwinia amylovora]. Ext. Bull. E, Coop. Ext. Serv. Mich. State Univ. (841) p.4. EL-81 Conway, K. E., L. S. Morrison, and P. W. Pratt. 1979. Fire blight of ornamentals and fruit trees. Okl. State Ext. Serv., Fact Sheet 7615. CANADA XII-A-99 Berg, C. W. and L. N. Gibbins. 1983. Impairment of substrate oxidation in the cytoplasm membrane of the fire-blight organism Erwinia amylovora. Can. Jour. Plant Path. 5:1-6. XII-A-100 Wall, R. E. 1983. Fire blight of wild raspberry on clear-cut forest areas. Can. For. Serv. Res. Notes 3:2-3. XII-A-101 Quamme, H. A. and G. A. Spearman. 1983. Cultivars 'Harvest Queen' and 'Harrow Delight' pear. HortScience 18:770-772. **NETHERLANDS** XII-F-75 Meijneke, C. A. R. 1983. Bacterievuur - de ontwikkeling van de situatie in 1982. Neth. Plant Prot. Serv. Yearb. (1982):69-76. Hoekstra, A., J. Cuijpers, P. Bolier and others. 1983. XII-F-76 Advieslijst waardplanten bacterievuur; alternatieven bacterievuurgevoelige gewassen. Landschap 16:25-27. XII-G-102 Billing, E. 1982. Fireblight. Jour. Royal Hortic. Soc. 108:206-210. XII-G-103 Billing, E., S. D. Trowell, M. M. Fuller, and others. 1982. Fireblight (Erwinia amylovora). Rep. East Malling Res. Sta. (1982):79-80.

XII-G-104

Anonymous. 1981.

Sta. (1981):102-109.

Diseases and pests of fruit. Rep. Long Ashton Res.

Blakeman, J. P. 1981. XII-G-105

The ecology of Erwinias on aerial plant surfaces, p.411-431. In Microbial ecology of the Phylloplane.

Academic. Press, London.

DENMARK

Middelsen, S. A. 1981. XII-H-40

Agricultural meteorology: Report from the soil Meteorology Project 1980. Tidsskrift for Planteavl.

85:73-74.

XII-H-41 Mosegaard, J. 1980.

Notes on fireblight through 200 years. Ugeskrift for

Jordbrug 125:1148-1150.

WEST GERMANY

XII-L-107 Hoppe, J. H. 1976.

Der Feuerbrand - eine Bedrohung für den Kernobstbau?

Hannov. Land - und Forstwirtsch. Zeitung. 129:26.

Persiel, F. 1979. XII-L-108

> Cotoneaster. Resistenz die Feuerbrand gegen

krankheit. Gaertnerboerse und Gartenwelt 79:628-629.

XII-L-109 Stuenz, E. 1979.

> Feuerbrand. Bauernblatt fuer Schleswig-Holstein.

33:80.

Schulz, F. A. 1981. XII-L-110

Biologischer Pflanzenschutz auch gegen

Krankheitserreger? Das Beispiel Feuerbrand. Deutsch.

Baumschule 33:519.

XII-L-111 Michel, H. G. 1981.

> Die Feuerbrandkrankheit gefaehrdet Obst und

Ziergehoelze. Deutsch. Gartenbau 35:1902-1904.

XII-L-112 Granhall, I. 1981.

> Alte und neue probleme der Pflanzenquarantaene.

Gesunde Pflanz. 33:122-123.

XII-L-113 Rapp, L. and J. Richter. 1982.

Uberprufung von desinfektionsmoglichkeiten beim arbeiten mit feuerbrand (Erwinia amylovora Burrill)

Winslow et al.). Nachrichtenbl. des Deutschen

Pflanzenschutzdienstes 34:177-178.

XII-L-114 Vorwohl, G. 1982.

Feuerbrandkrankheit und bienenhaltung. Allgemeine

Deutsch. Imkerzeitung 16:138-140.

XII-L-115 Feuerbrand D. 1982.

> Feuerbrand. Biol. Bundesanst. Land Forstwirtsch.

Merkbl. 0 (30):1-8.

- XII-L-116 Zeller, W. 1983.

 Resistenzprufung von Apfel und Birnensorten gegen den Feuerbrand. Mitt. des Sudtiroler Beratungsringes 20:162-163.
- XII-L-117 Zeller. W. 1983.

 Resistenzprufung von Apfel und Birnensorten gegen den Feuerbrand (<u>Erwinia amylovora</u>) in Schleswig-Holstein. Obstbau 8:266-268.
- XII-L-118 Zeller, W. 1983.

 Arbeitssitzung Feuerbrand v. 25.1.1983 im Pflanzenschutzamt Bonn-Bad Godesberg. Nachrichtenbl. Deut. Pflanzenschutzd. (Braunschweig) 35:140-141.
- XII-L-119 Zeller, W. 1983.
 Feuerbrand. Information zum integrierten
 Pflanzenschutz. Nachrichtenbl. Deut.
 Pflanzenschutzd. (Braunschweig) 35:174.
- XII-L-120 Zeller, W. 1983.

 Resistance of pome fruit varieties to fireblight

 (Erwinia amylovora) in the Fed. Rep. of Germany. Acta

 Hortic. 140:35-42.
- XII-L-121 Burlez, W. and W. Zeller. 1982.

 Feuerbrandbefall kann jetzt genau vorhergesagt werden. Taspo 116:7.
- XII-L-121 Sanftleben, U. and W. Zeller. 1983.

 Erster Computer-Einsatz zur Feuerbrandprognose im Baumschulbereich. Taspo Magazin 3:24-25.
- XII-L-122 Graf, H., Th. Kock, E. Lux-Wellenhof, J. Martin,
 S. Massfeller, and W. Zeller. 1983.
 Richtlinien zur Prufung von Bakteriziden gegen Erwinia
 amylovora (Feuerbrand) an Obst und Ziergeholzen. Biol.
 Bundesanst. Land. und Forstwirtschaft 12(1):1-6.

EAST GERMANY

- XII-M-15 Schaefer, H. J., W. Ficke, and H. Kleinhempel. 1982.

 Der Feuerbrand am Kernobst. Nachrichtenbl. für den Pflanzenschutz in der DDR 36:212-214.
- XII-M-16

 Fischer, C., H. J. Schaefer, M. Fischer, B. Dathe, and W. Ficke. 1983.

 Erste Ergebnisse der Resistenzzuchtung gegen Feuerbrand, Erwinia amylovora (Burrill) Winslow et al. bei Kernobst, 1. Mitt. Apfelsorten. Archiv für Zuchtungsforschung 13:137-146.
- XII-M-17 Fischer, M., C. Fischer, H. J. Schaeffer, and W. Ficke. 1983.

 Erste Ergebnisse der Resistenzzuchtung gegen Feuerbrand, Erwinia amylovora (Burrill) Winslow et al bei Kernobst, 2 Mitt. Apfel- und birnenunterlagen. Archiv für Zuchtungsforschung 13:183-191.

SWITZERLAND

XII-0-31 Moser, K. 1982.

Bericht uber die Fruhjahrskonferenz des vdsb fur die Sektions- und Kantonalprasidenten vom 2-7. Marz 1982 in Luzern. Schweiz. Bienen Zeitung 105:312-316.

XII-0-32 Anonymous. 1982.

Verordnung uber die Bekampfung der San-Jose Schildlaus, des Feuerbrandes und der gemeingefahrlichen Obstvirosen. Schweiz. Bundesrat, Artikel 60, April 28.

POLAND

XII-P-10 Sobiczewski, P. 1983.

Epidemiologia zarazy ogniowej. Ogrodnictwo 10:5-8.

XII-P-11 Sobiczewski, P. 1983.

Problemy ochrony drzew owocowych przed zaraza ogniowa. Ogrodnictwo 11:3-7.

FRANCE

XII-Q-58 Paulin, J. P. 1978.

Recente extension du feu bacterien en France. Jour. Fruit. d'Avignon-Montfavet 84:16-17.

XII-Q-59 Angiboust, A. 1979.

Choosing pear cultivars: earliness and fire blight resistance. France Agric. 35(1760):55-56.

XII-Q-60 Fieret, M. and M. Large. 1979.

Le feu bacterien dans le sud-ouest. Un danger pour le verger français. Phytoma (no.307):33-35.

XII-Q-61 Anonymous. 1979.

N'introduisons pas le feu bacterien dans nos cultures. Hortic. Franc. 103:29-30.

XII-Q-62 Anonymous. 1979.

Le feu bacterien menace les cultures. Pour repondre aux questions de vos clients sur le feu bacterien. Graines et Jardins 16:22-23.

XII-Q-63 Bassino, J. P. 1980.

Les problemes poses par la protection phytosanitaire des poiriers en France. Bull. Sect. Reg. Quest Palearctique 3(7):12-15.

XII-Q-64 Boue, H. 1980.

Situation du feu bacterien dans le Sud-Quest de la France. Sect. Reg. Quest Palearctique Bull. 3(7):42-44.

- XII-Q-65 Thibault, B. 1980.
 Aspects actuels de l'amelioration du poirier pour la resistance au feu bacterin. Fruit Belge 48(390):123-129.
- XII-Q-66 Thibault, B. 1980.
 Aspects actuels de l'amelioration du poirier pour la resistance au feu bacterien. Arboriculture Fruitiere 316:59-61.
- XII-Q-67 Lancrenon, P. 1980.

 Poiriers, pommiers: alerte au feu bacterien.

 Agri-Sept 777:28-29.
- XII-Q-68 Mathys, G. 1981.

 Einige gedanken zur pflanzenquarantaene. Gesunde Pflanz. 33:49-53.
- XII-Q-69 Pinault, M. 1981.

 Peut-on vivre avec le feu bacterien? France Agricole 36:33-37.
- XII-Q-70 Large, M. 1981.

 Le feu bacterien en France (Erwinia amylovora).

 Situation actuelle et perspectives. Bull. Tech.
 d'Information 357:93-109.
- Fougeroux, A. 1981.

 La situation du feu bacterien en France a la fin de 1980. Phytoma 329:18-19.
- XII-Q-72 Carles, L. 1983. Le feu bacterien. Arboric. Fruit. 30:348:50-53.
- XII-Q-73 Anonymous 1983.

 Mesure de lutte contre le feu bacterien des rosacees.

 Def. Veg. INRA-Pathol. 37:(219):44-51.
- XII-Q-74 Anonymous 1983.

 La lutte contre le feu bacterien. Phytoma 347:41-42.
- XII-Q-75 Anonymous 1983.

 Selection pour la resistance au feu bacterien.
 Agronomie 3(1):90.
- XII-Q-76
 Paulin, J. P. 1983.
 Choix d'un isolat d'<u>Erwinia amylovora</u> pour la determination de la sensibilite varietale au feu bacterien. Bull. Sect. Reg. Quest Palearc. 4:171-177.
- XII-Q-77 Acquart-Romon, C. J. P. Paulin, and D. Payen. 1983.
 Aspects climatiques de l'epidemiologie du feu bacterien en France. C. R. Acad. Agric. de France p.944-955.
- XII-Q-78 Thibault B., C. Welcker, and Y. Lespinasse. 1983.

 Heritability of the character "secondary blooming" on pear (Pyrus communis). Acta Hortic. 139:181-193.

Anonymous. 1983. XII-Q-79 Liste des plantes proposees par la FNPHP et le CNIH en remplacement des plantes tres sensibles au feu bacterien. L'Hortic. Franc. (Decembre):9-12. XII-Q-80 Mathys. G. 1983. Rapport du groupe de travail pour l'etude problemes phytosanitaires specifiques aux cultures mediterraneenes. Europ. Mediter. Plant Prot. Org. Publ. 81, 34pp. XII-Q-81 European and Mediterranean Plant Protection Organisation. 1983. New infection by <u>Erwinia amylovora</u> detected in the departement of <u>Loiret</u>; infection at St. Julien-en-Genevois (Haute Savoie). Europ. & Mediter. Plant Prot. Org. Rept. RSE 448, 1 p. European and Mediterranean Plant Protection XII-Q-82 Organisation. 1983. Erwinia amylovora (Burrill 1882) Winsl. et al. 1920. Europ. & Mediter. Plant Prot. Org. Bull. 13(1):1-14. XII-Q-83 European and Mediterranean Plant Protection Organisation. 1981. Data sheets on quarantine organisms. EPPO list A2. Europ. & Mediter. Plant Prot. Org., Mimeogr. material. BELGIUM XII-R-18 Laroche, M. and M. Verhoyen. 1981. Serological identification of phyto bacteria in Belgium. 33rd Intern. Symp. on Crop Prot., Pt. 3 46:757-772. XII-R-19 Laroche, M. and M. Verhoyen. 1982. Serological identification of Erwinia amylovora using immuno diffusion. 34th Intern. Symp. Crop Prot., Pt. 3 47:1083-1094. XII-R-20 van Vaerenbergh, J. 1982. Le feu bacterien des rosacees. Fruit Belge 50(398):142-152 XII-R-21 Porreye, W. 1982. Le feu bacterien, situation actuelle, perspectives et methodes de lutte. Fruit Belge 50(398):152-155.

> de Greef, M., O. van Laere, and L. de. Wael. 1982. L'apiculture et le feu bacterien en arboriculture et en culture ornementale. Agricontact 130:4.

XII-R-22

XII-R-23 Destinay, P. 1982. Tableau d'identification des aubepines. Environment 6:1-7. XII-R-24 Mergaert, J., L. Verdonck, J. Swings, K. Kesters, and J. Deley. 1982. The use of API systems for the identification of Erwinia species. Med. Fac. Landbouww. Rijksuniv Gent. 47/3:1095-1100. XII-R-25 De Greef M., O. Van Laere, and L. De Wael. 1982. Bacterievuur en bijenteelt. Maandblad Vlaamse Imkersbond 68 (4):126-129. XII-R-26 Vantomme, R., J. Swings, and J. de. Ley. 1983 Le feu bacterien, une menace permanente. Rev. de L'Agric. 36:257-262. XII-R-27 Porreye, W., and T. Deckers. 1982 lst steps in chemical control of <u>Erwinia</u> amylovora. 34th Intern. Symp. Crop Prot. Pt. 3 47:1101-1104 XII-R-28 Deckers, T., and W. Porreye. 1983 Ervaringen met Copac e als bestrijdingsmiddel tegen bacterievuur. Fruitteelt 73:156-157. XII-R-29 Porreye, W., and T. Deckers. 1983 Methodes de prevention de developpement du feu bacterien (<u>Erwinia amylovora</u>). Parasitica 39;no 2. XII-R-30 Deckers, T., and W. Porreye. 1983. La lutte contre le feu bacterien, Erwinia amylovora (Burr.) Winslow et al. Fruit Belge 51(402):164-167. LUXEMBURG XII-S-1 European and Mediterranean Plant Protection Organisation. 1983. New infection by fire blight detected in Luxemburg. Europ. & Medit. Plant Prot. Org. Rept. 499:1. ITALY Anonymous. 1981. XII-T-23 Colpo di fuoco batterico delle pomacee. Giornale di Agricoltura 91:36-37. XII-T-24 Tanga, R. M., and A. de Bono. 1982. Il colpo di fuoco. Lazio Agricolo 4:24-27 Agnolin, C., R. De. Clauser, and S. Emer. 1982. XII-T-25 Abbiamo visto da vicino colpo di fuoco batterico (viaggio di studio in Germania). Terra Trentina

28:25-30.

XII-T-26 Calzolari, A., U. Mazzucchi, and C. Gasperini. 1982.

Cross-reactions between Erwinia amylovora and other bacteria in immunofluorescence staining using different antisera. Phytopath. Mediter. 21:110-112

XII-T-27 Italian Ministry of Agriculture. 1983.

Disposizioni fitosanitarie per l'importazione di piante ospiti di Erwinia amylovora. Decr. Min., Gaz. Uff. Rep. Ital. 1:6-7

SPAIN

Palazon Espanol, J., A. Alfaro Moreno, and M. C. Rodriguez. 1974.

El fuego bacteriano (<u>Erwinia amylovora</u>) de las rosaceas: Un grave peligro para nuestra fruticultura. Inst. Nac. Invest. Agrarias, Madrio (Spain) 46pp.

XII-U-5 Sampayo, M., and I. J. Palazon. 1980.

Medidas que se estan tomando en Espana respecto al fuego bacteriano de las rosaceas (Erwinia amylovora Burril). Org. Intern. de Lutte Biolog. Sect. Reg. Quest Palearc. 3:45-49.

XII-U-6 Sampayo Fernandez, M. 1979.

El fuego bacteriano (una enfermedad bacteriana producida por <u>Erwinia amylovora</u>). Agricultura 48:697-700.

XII-U-7

Arias Firalda, A., M. Blasco Escudera, and others
1982.

El Fuego Bacteriano de las Rosaceas. II. Min. de
Agric., Pesca y Aliment., Booklet, 25pp. illus. with
36 slides.

YUGOSLAVIA

XII-W-l Kispatic, J. 1975.

Pales cvijeta Kruske-nova opasnost za jezgricave vocke. (Pear flower fire-blight new danger for temperate fruit trees). Biljna Zastita 19:133-137.

RUSSIA

Mikhailichenko, N. V., N. I. Kochetova, and Y. V. Kochetov. 1980.

Cell composition of different types of bacterial colonies from the amylovorous group (Erwinia amylovora blight of fruit). Soviet Agriculture Sciences (USA) 8:24-27.

XIII-A-32 Kalinichenko, R. I. and L. G. Panfilova. 1980.

Express diagnosis of the pathogen of bacterial fire blight of fruit crops. Selskoe Khoz. za Rubezhom 12:27-29.

XIII-A-33 Kalinichenko, G. V. and R. I. Kalinichenko. 1983. Characteristics of fruiting of pear under bacterial infection in the crimea. Sel'Skokhoz. Biologiya 4:68-71.

EGYPT

XIII-E-7 Barakat, F. M. and N. A. Elsa. 1977. A laboratory method for testing virulence in Erwinia amylovora strains. Annals Agric. Sci.

Moshtohor 8:167-176.

XIII-E-8 Barakat, F. M. 1979.

Variations in disease expression and multiplicity rate in host tissues, induced by different mutant strains of <u>Erwinia amylovora</u>. Proc. 3rd Egypt. Phytopath. Congr., p. 299-310.

VENEZUELA

XIII-U-1 Zambito Leo, L., and F. L. Lukezic. 1980.

Efecto de periodo de exposicion de luz sobre la produccion de isoflavonoides en hojas (Glycine max) infiltradas con Erwinia amylovora y agua. Jornadas Intern. Escuela Agron., Univ. Centro Occid. Lisandro Alvarado, Barquisimeto, p. 50-51.



LIST OF PERSONS INTERESTED IN FIRE BLIGHT $\frac{1}{2}$

Abdel-Rahman, M., Fertilizer-Chemical Division, Agway Inc., P.O. 4933, Syracuse, New York 13221. (315-477-6176)	(1)	USA
Abo-El-Dahab, M. K., Plant Pathology Dept., Faculty of Agric., Univ. of Alexandria, Alexandria, Egypt (71960)	(1)	EGY
Agriculture Canada, Library, Records Div., Ottawa, Ontario KIA OC5, Canada	(2)	CND
Aldwinckle, H. S., Department of Plant Pathology, N.Y. State Agric. Expt. Station, Geneva, New York 14456. (315–787–2317)	(1)	USA
Alston, F. H., Fruit Breeding Department, East Malling Research Station, East Malling, Maidstone, Kent, ME19 6BJ, England. (0732–843833)	(1)	UK
Andersen, H., The Government Plant Protection Service, Gersonsvej 13, 2900 Hellerup, Denmark. (01–620787)	(1)	DK
Ark, P. A., St. Pauls Towers, 100 Bay Place, Apt. 1915, Oakland, California 94610. (415-835-4700, ext. 298)	(4)	USA
Arsenijevic, M., Faculty of Agriculture, Institute for Plant Protec., V. Vlakovica 2, 21000 Novi Sad, Yugoslavia. (021-58-366)	(3)	YUG
Balavoine, P., Service de la Protection des Vegetaux, Cite Administr., 59048 Lilla, France.	(2)	FR
Barrat, J. G., West Va. University Expt. Farm, Kearneysville, West Virginia 25430. (304-876-6353)	(1)	USA
Bates, J. J., Biological Research Center, Imperial Chemicals Inc., P.O. Box 208, Goldsboro, North Carolina 27530. (919-736-3030)	(2)	USA
Baumm, L. H., Institute fur Angewandte Botanik, Univ. of Hamburg, Marseillerstrasse 7, 2000 Hamburg 36, West Germany (040/5 11.89.12)	(1)	BRD

Names underlined are contact persons for preparation of fire blight newsletter. Numbers in parentheses following addresses are local telephone numbers, and those in column at right indicate activity or interest in fire blight:

Actively engaged in fire blight research;

Indirectly interested in fire blight;

^{3.} Interested in fire blight, but located in region where disease is not present:

^{4.} Retired but still interested in fire blight activities.

Baykal, N., Agric. Univ. Ziraat Fakultesi, Fitopatoloji Kursusu, Ankara, Turkey.	(3)	TUR
<pre>Bazzi, C., Laboratorio Fitobatteriologia, Istituto Patol. Vegetale, via Filippo Re 8, 40126 Bologna, Italy. (051-236175)</pre>	(3)	ITA
Beer, S. V., Department of Plant Pathology, Cornell University, Ithaca, New York 14853. (607-256-3259)	(1)	USA
Bell, R. L., U.S. Department of Agriculture, Appalachian Fruit Research Station, Rt. 2, Box 45, Kearneysville, West Virginia 25430. (304-725-3451, ext. 21)	(1)	USA
Benjama, A., Laboratoire de Phytiatrie et Phytobacteriologie, Institut Nationale de la Recherche Agronomique, B.P. 415, Rabat, Morocco.	(3)	MOR
Bennett, R. A., Redditch College, Peakman Street, Redditch, Worcester B98 8DW, England	(2)	UK
Bergna, D. A., Estacion Experimental Alto Valle, Casilla de Correo 52, 8332 General Roco, Rio Negro, Argentina. (0941-22248)	(3)	ARG
Berry, D. W., Jackson County Extension Office, 1301 Maple Grove Drive, Medford, Oregon 97501.	(2)	USA
Beutel, J. A., Department of Pomology, University of California, Davis, Calif. 95616. (916–752–0507)	(1)	USA
Biehn, W., Ciba Geigy Corporation, R + D Agric. Div., Box 11422, Greensboro, North Carolina 27409.	(2)	USA
Biggs, A. R., Agriculture Canada, Research Station, Vineland, Ontario LOR 2EO (416-562-4113)	(2)	CND
Billing, Eve, 4 Fromandez Drive, Horsmonden, Tonbridge, Kent TN12 8LN, England (089-272-2807)	(1)	UK
Bolay, A., Section de Phytopathologie, Station Federale de Recherches Agronomiques de Changins, 1260 Nyon, Switzerland. (022-615451)	(3)	SWT
Bonn, W. G., Agriculture Canada, Research Station, Harrow, Ontario NOR 1GO, Canada. (519-738-2251)	(1)	CND
Bouma, S., Research Station for Arboriculture, P.O. Box 118, 2770 AC Boskoop, The Netherlands. (01727-3220)	(1)	NL
Bredemeier, D., Universidade Federal de Santa Maria, Departamento de Fitotecnia, 97-100 Santa Maria, Rio Grande do Sul, Brazil.	(3)	BRA
Brulez, W., Biologische Bundesanstalt, Institut fur Pflanzenschutz, Schlosskoppelweg 8, 2305 Heikendorf, West Germany. (0431–23495).	(1)	BRD

Burkowicz, A., Institut Sadownictwa, 83-111 Milobadz, Poland.	(1)	POL
Burr, T. J., Department of Plant Pathology, N.Y. State Agric. Expt. Station, Geneva, New York 14456. (315-787-2312)	(2)	USA
Bushong, J. W., Agrichemicals Div., 3M Center, Bldg 223-IN-05, 3M Company, St. Paul, Minnesota 55144 (612-736-0930)	(2)	USA
Button, J., Box 86, Ceres 6835, Republic of South Africa.	(3)	SA
Byrde, R. J. W., Long Ashton Research Station, Bristol BS18 9AF, England. (027-580 2181)	(1)	UK
Callu, D., Service de la Protection des Vegetaux, Cite Administr., 67000 Strasbourg, France	(2)	FR
Calzolari, Alessandra, Osservatorio per le Malattie delle Piante, Via di Corticella 133, 40129 Bologna, Italy. (051-352917)	(3)	ITA
Cameron, H. R., Department of Botany & Plant Pathology, Oregon State University, Corvallis, Oregon 97330. (503-754-4044)	(2)	USA
<u>Cao</u> , R., Department of Plant Protection, Zhejiang Agricultural University, Hangzhou, Zhejiang, Peoples Republic of China. (42605)	(3)	CHI
Carlson, R. F., Department of Horticulture, Michigan State University, East Lansing, Michigan 48823. (517–355–5200)	(2)	USA
Carroll, V. J., Chemicals Division, Pfizer Inc., 235 East 42nd Street, New York, New York 10017. (212-573-2643)	(1)	USA
Cartwright, D. N., Plant Quarantine Div., South AustrDept. of Agric., Box 1671, G.P.O., Adelaide, South Australia 5001. (08-2660911)	(3)	AUS
Cazelles, O., Station Federale de Recherches Agronomique de Changins, 1260 Nyon, Switzerland.	(3)	SWT
Centre for Agricultural Publishing and Documentation, P.O. Box 4, 6700 AA Wageningen, The Netherlands.	(2)	NL
Chandler, D., 1006 S. 32nd Avenue, Yakima, Washington 98902. (509-253-3414)	(2)	USA
Chouibani, M., D.P.V.C.T.R.F., B. P. 1308, Rabat, Morocco	(3)	MOR
Christensen, F. G., The Royal Veterinary and Agricultural University, Arboretum, 2970 Horsholm, Denmark. (02–860641)	(2)	DK

Chronica Horticulturae (Editor), Geertjesweg 106, 6706 EE Wageningen, The Netherlands.	(2)	NL
Civerolo, E. L., Fruit Laboratory, U.S. Department of Agriculture, Room 111, Building 004, BARC-West, Beltsville, Maryland 20705. (301–344–4651)	(2)	USA
Clayton, C. N., Department of Plant Pathology, North Carolina State University, Raleigh, NC 27607. (919–737–2721)	(4)	USA
Cline, R. A., Horticulture Research Institute of Ontario, Vineland Station, Ontario LOR 2EO, Canada. (416–562–4141)	(2)	CND
Conceicao, J., Direccao General de Proteccao da Producao Agricola, Tapada da Ajuda, 1300 Lisbon, Portugal.	(3)	POR
Cooper, R. M., School of Biology Sciences, University of Bath, Claverton Down, Bath, Avon BA2 7AY, England	(1)	UK
Cornils, H., Inst. fur Angewandte Botanik, Univ. of Hamburg, Marseillerstr. 7, 2000 Hamburg 36, West Germany. (040–4123–2359)	(1)	BRD
Coulombe, L. J., Agriculture Canada, P.O. Box 457, St. Jean, Quebec J3B 6B8, Canada. (514–346–4494)	(2)	CND
Covey, R. P., Tree Fruit Research Center, 1100 North Western Avenue, Wenatchee, Washington 98801. (509-663-8181)	(1)	USA
Crassweller, R. M., Cooperative Extension Service, University of Georgia, Athens, Georgia 30602. (404-542-2861)	(2)	USA
Crowe, A. D., Agriculture Canada, Tree Fruit Section, Research Station, Kentville, Nova Scotia B4N 135, Canada. (902-678-2171)	(2)	CND
Cummins, J. N., Department of Pomology & Viticulture, N.Y. State Agr. Expt. Station, Geneva, New York 14456. (315-787-2233)	(1)	USA
Dale, T., Norwegian Plant Inspection Service, P.O. Box 94, Okern, Oslo 5, Norway. (02-224760)	(3)	NOR
Davidson, J. G. N., Agriculture Canada, Research Station, Box 29, Beaverlodge, Alberta TOH OCO, Canada. (403-354-2212)	(2)	CND
Davidson, S. H., 408 Troy Avenue (Woodcrest), Wilmington, Delaware 19804. (302-994-1875)	(4)	USA
Deckers, T., Opzoekingsstation van Gorsem, Brede Akker 3, 3800 St. Truiden, Belgium. (Oll-682019)	(1)	BLG

De Ley, J., Lab. voor Microb. en Microb. Genetica, Rijksuniv. Gent, K. L. Ledeganckstr. 35, 9000 Gent, Belgium. (22–78–21)	(1)	BLG
<u>Dinesen</u> , A., Botany Department, State Plant Pathology Institue, Lottenborgvej 2, 2800 Lyngby, Denmark. (0287–2510)	(2)	DK
Dobra, A., Catedra de Fitopatologia, Facultad de Ciencias Agrarias, Universidad Nacional del Comahue, 8303 Cinco Saltos, Argentina.	(3)	ARG
<u>Douglas</u> , S. M., Dept. of Plant Pathology, Conn.Agric. Expt. Sta., New Haven, Conn. 06504 (203-789-7222)	(2)	USA
<pre>Drake, C. R., Department of Plant Pathology & Physiology, Virginia Polytechnic Institute, Blacksburg, VA 24061. (703-961-5251)</pre>	(2)	USA
Duben, J., Bayer AG, Pflanzenschutzberatung, 5090 Leverkusen, West Germany.	(2)	BRD
Dye, D. W., Plant Diseases Division, Dept. of Scientific & Industr. Research, Private Bag, Auckland, New Zealand. (893660)	(2)	NZ
Egolf, D. R., U. S. National Arboretum, 3501 New York Ave., Washington, D. C. 20002. (202-472-9277)	(1)	USA
Egli, T., Ciba-Geigy Chem. Company Ltd., AC 2.82, 4002 Basel, Switzerland.	(3)	SWT
Ellis, M. A., Department of Plant Pathology, Ohio Agric. Research and Devel. Center, Wooster, Ohio 44691. (216-263-3700)	(1)	USA
Ercolani, G. L., Istituto di Microbiologia Agraria e Tecnica, Facolta di Agraria, Via Amemdola 165/A, 70126 Bari, Italy. (080–339422)	(3)	ITA
Erskine, J. M., Inst. of Natural Resources, Univ. of Natal, P.O. Box 375, Pietermaritzburg 3200, South Africa. (0331–21344)	(3)	SA
Evans, I. R., Plant Pathology Laboratory, 605 Agriculture Bldg., 9718-107 St., Edmonton, Alberta T5K 2C8, Canada. (403-427-5350)	(2)	CND
Feliciano, Ascunia J. (Connie), EMBRAPA/UEPAE de Cascata, Caixa Postal 403, Pelotas 96.100, Rio Grande do Sul, Brazil.	(3)	BRA
Fideohelli, C., Istituto Sperimentale per la Frutticoltura, Via di Fioranello n. 52, Ciampino Aeroporto, 00040 Rome, Italy.	(3)	ITA

Fox, R. T. V., I.C.I., Plant Protect. Division, Jealott's Hill Res. Station, Bracknell, Berkshire RG12 6EY England. (0344–24701)	(2)	UK
Franz, W., Amt fur Land-und Wasserwirtschaft, Abt. Pflanzensch., Schonbockener Str. 102, 2400 Lubeck, West Germany. (0451-45551)	(2)	BRD
French, J. R., FMC Corp., 100 Niagara Street, Middleport, New York 14105. (716-735-3761, ext. 361)	(2)	USA
Fucikovsky, L., Centro de Fitopatologia, Colegio de Postgraduados, Escuela Nacional de Agricultura, A.P.#85, 56230 Chapingo, Mexico. (5-85-45-55, ext. 5556)	(2)	MEX
Gantotti, B. V., Department of Bacteriology, University of California, Davis, California 95616. (916-756-0283).	(1)	USA
Garibaldi, A., Istituto di Patologia Vegetale, Via Giuria 15, 10126 Torino, Italy. (011-6505236)	(3)	ITA
Garrett, Connie M. E., Dept. of Plant Pathology, East Malling Research Station, Maidstone, Kent, ME19 6BJ, England (0732-843833)	(1)	UK
Gates, D., Agric. Chemicals Div., 3-M Company, 223-1 N.E., St. Paul, Minnesota 55144. (612-736-9476)	(2)	USA
Geenen, J., Rijksstation voor Plantenziekten, Burg. van Gansberghelaan 96, 9220 Merelbeke, Belgium.	(2)	BLG
Gibbins, L. N., Department of Microbiology, University of Guelph, Guelph, Ontario NIG 2Wl, Canada. (519–824–4120, ext. 3477)	(2)	CND
Goodman, R. N., Dept. of Plant Pathology, University of Missouri, Columbia, Missouri 65211. (314-882-7043)	(1)	USA
Goto, M., Laboratory of Plant Pathology, Faculty of Agriculture, Shizuoka University, 836 Ohya, Shizuoka 422, Japan. (0542–37–1111, ext. 827)	(3)	JAP
Graberg, M., National Board of Agriculture, Plant Protection Service, 551 83 Jonkoping, Sweden. (036-16.94.20)	(3)	SWD
Graf, H., Obstbauversuchsanstalt, Westerminnerweg 22, 2155 Jork, West Germany. (04162-7511)	(2)	BRD
Grimm, R., Federal Res. Station for Fruit-growing, Viticulture and Hortic., 8820 Wadenswill, Switzerland. (01-780.13.33)	(3)	SWT
Gupta, G. K., Fruit Pathology Laboratory, Regional Fruit Res. Station, Black Rock, Mashobra, Simla 7, H.P., India. (8261).	(3)	IND

Gwynne, D. C., Agricultural Development and Advisory Service, Min. of Agric., Fisheries, and Food, Burghill, Rd., Westbury-on-Trym, Bristol BS10 6NJ, England. (0272–500000)	(1)	UK
Hale, C. N., Plant Diseases Div., Dept. of Scientific and Industrial Research, Private Bag, Auckland, New Zealand (893660)	(1)	NZ
Harnish, W., Agric. Chem. Div., Food & Machinery Corporation, 100 Niagara Street, Middleport, New York 14105. (716–735–3761)	(2)	USA
Heimann, Mary Francis, Dept. Plant Pathology, University of Wisconsin, Russell Labs, 1630 Linden Drive, Madison, Wisconsin 53706. (608–262–1426)	(1)	USA
Heybroek, H. M., Dorschkamp Research Inst. for Forestry and Landscape Planning, P. O. Box 23, 6700 AA Wageningen, The Netherlands. (08370—19050)	(1)	NL
Hickey, K. D., Fruit Research Laboratory, Penn. State Univ., Box 309, Biglerville, Pennsylvania 17307. (717–677–6116)	(2)	USA
Hignett, R. C., Dept. of Plant Pathology, East Malling Research Station, Maidstone, Kent ME19 6BJ, England	(1)	UK
Hildebrand, E. M., 11092 Timberline Drive, Sun City, Arizona 85351. (602-977-5326)	(4)	USA
Hoppe, H., Pflanzenschutzamt Hannover, Bez. Stelle Bremervorde, Neue Str. 22, 2140 Bremervorde, West Germany.	(2)	BRD
Horricks, J., Alberta Agriculture, 9718 107 Street, Edmonton, Alberta T5K 2C8, Canada. (403-427-5350)	(2)	CND
Howard, R. J., Alberta Hort. Res. Center, BAG Service 200, Brooks, Alberta TOJ OJO, Canada. (403-362-3391)	(2)	CND
Hunter, C. L., Soils and Crops Branch, Ontario Ministry of Agriculture and Food, P. O. Box 587, Simcoe, Ontario N3Y 4N5, Canada. (519–426–7120)	(1)	CND
<pre>Isenbeck, Margot, Institut fur Phytopathologie, Universitat Kiel, Olshausenstr. 40-60, 2300 Kiel, West Germany. (0431-880-2996)</pre>	(1)	BRD
<u>Janick</u> , J., Department of Horticulture, Purdue University, West Lafayette, Indiana 47907. (317-494-1329).	(1)	USA
Johnson, D. E., 3310 Jefferson Avenue, Yakima, Washington 98902.	(2)	USA
Jones, A. L., Department of Botany & Plant Pathology, Michigan State University, East Lansing, Michigan 48823. (517–355–4573)	(2)	USA

Jorgensen, H. A., National Plant Pathology Institute, Lottenborgvej 2, 2800 Lyngby, Denmark. (01-8725-10)	(2)	DK
Joseph, E., Service Phytosanitaire, Div. de l'Agriculture, Hartenhofstr. 5, 3003 Bern, Switzerland.	(3)	SWT
Kado, C. I., Department of Plant Pathology, University of California, Davis, California 95616. (916-752-0325)	(2)	USA
Kappel, F. Department of Horticulture, University of Guelph, Guelph, Ontario NIG 2W1, Canada. (519-824-4120)	(1)	CND
Kato, T., Research Department - Pesticides Div., Institute for Biological Science, Sumitomo Chemical Co., Ltd., 4-2-1, Takatsukasa, Takarazuka, Hyogo, 665, Japan.	(3)	JAP
Kleinhempel, H., Inst. fur Phytopathologie, Akad. Landwirtsch. Wissensch., Theodor-Roemer-Weg 4, 4320 Aschersleben, East Germany. (5141)	(3)	DOR
Klement, Z., Dept. of Pathophysiology & Disease Resistance, Plant Protection Institute, Herman Otto u. 15, P.O. 102, 1525 Budapest, Hungary. (358-137)	(3)	HUN
Klos, E. J., Department of Botany & Plant Pathology, Michigan State Univ., East Lansing, Michigan 43823. (517–355–4680)	(1)	USA
<pre>Knosel, D., Inst. fur Angewandte Botanik, Univ. of Hamburg, Marseiller Str. 7, 2000 Hamburg 36, West Germany. (040-4123-2353)</pre>	(1)	BRD
Koenigshof, R., Pear Research Association, Box 4050, Kerlikowske Rd., Coloma, Michigan 49038. (616-849-2375)	(2)	USA
Kooistra, T., Plant Protection Service, Geertjesweg 15, P. O. Box 9102, 6706 EA Wageningen, The Netherlands. (08370-19001)	(1)	NL
Kraus, P., Bayerwerk, Pflanzenschutz Anwendungstechnik, Biologische Forschung, 5090 Leverkusen, West Germany. (02172-306081)	(3)	BRD
Kristensen, H. R., State Plant Pathology Institute, Lottenborgvej 2, 2800 Lyngby, Denmark	(2)	DK
Kroeker, G., Swedish Univ. of Agric. Sciences, Box 7036, 75007, Uppsala 7, Sweden. (013-102000)	(3)	CWZ
Kuc, J., Dept. of Plant Pathology, S-305 Agric. Sci. Center North, Univ. of Kentucky, Lexington, Kentucky 40506. (606-258-4978)	(2)	USA

Kudela, V., Institute of Plant Protection, Research Inst. of Plant Production, Drnovska 507, 16106 Prague 6 (Ruzyne), Czechoslovakia.	(3)	CZE
Kuhne, H., Pflanzenschutzamt der Ldw. Kammer Weser-Ems, Mars-la-Tour-Str. 9/11, 2900 Oldenburg, West Germany.	(2)	BRD
Kyle, Nancy E., 2222 N. Richland, Phoenix, Arizona 85006. (602–252–3060)	(1)	USA
Lacy, G. H., Department of Plant Pathology, Conn. Agric. Expt. Station, New Haven, Connecticut 06504. (203–789–7222)	(2)	USA
Laere, O. van, Research Station for Nematology and Entomology, Burg. van Gansberghelaan 96, 9220 Merelbeke, Belgium. (091–52.20.85)	(1)	BLG
Lamb, R. C., Department of Pomology & Viticulture, N. Y. State Agr. Expt. Station, Geneva, New York 14456. (315-787-2235)	(1)	USA
Landis, W. R., Agric. Chem. Development, MSD Agvet Division, P. O. Box 2000, Rahway, New Jersey 07065. (201-574-6605)	(2)	USA
Lane, D., Agriculture Canada, Research Branch, Res. Station, Summerland, British Columbia VOH 1ZO, Canada. (604–494–0401)	(2)	CND
Langeslag, J. J., Plant Protection Service, Geertjesweg 15, P. O. Box 9102, 6700 HC Wageningen, The Netherlands.	(2)	NL
Large, M., Service de la Protection des Vegetaux, Chemin d'Artigues B.P.47, 33150 Cenon La Morlette, France. (56-86.22.75)	(1)	FR
Larcoche, M., Centre d'Etudes des Phytobacterioses, Lab. de Phytopathologie, 3 Place Croix du Sud. Sci. 15 D, 1348 Louvain-La-Neuve, Belguim (010-41.81.81 ext. 3746)	(1)	BLG
Lecomte, P., Laboratoire Feu Bacterien, Lycee Agricole d'Oereluy, St. Paul, 40990 Les Dax, France. (58–74.11.93, ext. 22)	(1)	FR
Lehmann-Danzinger, H., Inst. fur Pflanzenpath. und Pflanzensch., Grisebachstr. 6, 3400 Gottingen, West Germ.	(2)	BRD
Lelliott, R. A., Agricultural Science Service, Harpenden Laboratory, Hatching Green, Harpenden, Herts AL5 2BD, England. (Harpenden 5241)	(2)	UK
Letal, J., Regional Crops Laboratory, Box 10, Olds, Alberta TOM IPO, Canada. (403–556–8421)	(2)	CND

Lombard, P. B., Department of Horticulture, Oregon State University, Corvallis, Oregon 97331. (503-754-3695)	(2)	USA
Lopez Gonzalez, M., Dept. Proteccion Vegetal, I.N.I.A., CRIDA 07, Moncada-Valencia, Spain. (739-1000)	(3)	SPN
Luchene, K., van, Ministerie van Landbouw, Dienst Planten- bescherming, Gebrs. Vandeveldestraat 68, 9000 Gent, Belgium.	(1)	BLG
Luepschen, N. S., Tree Disease Consulting Service, Rim Rock Campground, 73179 State Highway 64, Meeker, Color. 81641. (303–878–4486)	(2)	USA
Lux-Wellenhof, E., Ciba-Geigy GmbH, Postfach 11.03.53, 6000 Frankfurt 11, West Germany (0611-7155.257)	(2)	BRD
Maas Geesteranus, H. P., Research Institute for Plant Protection, Binnenhaven 12, P. O. Box 9060, 6700 GW Wageningen, The Netherlands. (08370-19151)	(1)	NL
Mansergas, A. J. F., Ministerio de Agricultura, Departamento de Fruticultura, Apartado 202, Zaragoza, Spain. (976-29 72 07)	(3)	SPN
Mappes, D., BASF, Fruit Experimental Station, P.O. Box 220, 6703 Limburgerhof, West Germany	(2)	BRD
Maroquin, C., Station de Phytopathologie de l'Etat, 13 Ave. Marechal Juin, 5800 Gembloux, Belgium.	(2)	BLG
Martins, J. M. S., Dept. Fitopatologia, Estacao Agronomica Nacional, 2780 Oeiras, Portugal.	(3)	POR
Massfeller, D., Pflanzenschutzamt der Ldw. Kammer Rheinland, Ludwig Erhard Str. 99, 5300 Bonn-2, West Germany. (02221-376931)	(1)	BRD
Mathys, G., European and Mediter. Plant Protect. Organ., l rue Le Notre, 75016 Paris, France. (870-77-94)	(2)	FR
Matthee, F. N., Department of Plant Pathology, Univ. of Western Cape, Bellville, South Africa.	(3)	SA
Mazzucchi, U., Laboratorio Fitobatteriologia, Istituto Patol. Vegetale, via Filippo Re 8, 40126 Bologna, Italy. (227401)	(3)	ITA
McPhee, R., Agriculture Canada, Research Station, Summerland, British Columbia VOH 1ZO, Canada. (604-494-7711)	(2)	CND
McSwan, I. C., Extension Plant Pathology, 1089 Cordley Hall, Oregon State University, Corvallis, Oregon 97331. (503–754–3472)	(4)	USA

Meijneke, C. A. R., Plant Protection Service, Geertjesweg 15, P. O. Box 9102, 6700 HC Wageningen, The Netherlands. (08370—19001)	(2)	NL
Mendoza H., A., Centro de Fitopatologia, Colegio de Postgraduados, 56230 Chapingo, Est. de Mexico, Mexico (5-85-45-55 ext. 5406)	(1)	MEX
Meyer, F. C., Catedra de Fitopatologia, Facultad de Ciencias Agrarias, Universidad Nacional del Comahue, 8303 Cinco Altos, Argentina.	(3)	ARG
Meyer, J., Amt fur Land-und Wasserwirtschaft, Abteilung Pflanzenschutz, Herzog-Adolf Strasse 1b, 225 Husum, West Germany. (04841–2746)	(1)	BRD
Michel, H. G., Landesanstalt fur Pflanzenschutz, Reinsburgstr. 107, 7000 Stuttgart - 1, West Germany. (0711/6676-2575 or 73)	(3)	BRD
Mickail, K. Y., Plant Pathology Research Inst., Agric. Research Centre, Cairo (Giza), Egypt (723000)	(1)	EGY
Miller, H. J., Plant Protection Service, Geertjesweg 15, P. O. Box 9102, 6700 HC Wageningen, The Netherlands.	(1)	NL
Miller, R. W., Dept. of Plant Path. and Physiol., Clemson Univ., Clemson, South Carolina 29631. (803-656-2335)	(2)	USA
Morehead, G. W., Farm Advisors Office, 4145 Branch Center Road, Sacramento, California 95827. (916–366–2013)	(1)	USA
Morton, H. V., Ciba-Geigy Corp., P.O. Box 18300, Greensboro, North Carolina 27419. (919-292-7100)	(2)	USA
Mosegaard, J., Dansk Plantekoleejer Forening, Elmedals Allee 33, 5250 Fruens Boge, Denmark.	(2)	DK
Muir, J., Alberta Agriculture, Research Station, Fairview, Alberta, TOH 1LO, Canada.	(2)	CND
Muller, H. J., Institut fur Phytopathologie, Theodor-Roemer Weg 1-4, 432 Aschersleben, East Germany.	(3)	DDR
Muller, K., Institut fur Pflanzenschutz der Ldw. Kammer Westfalen-Lippe, Kanalstr. 240, 4400 Munster, West Germany.	(2)	BRD
Norelli, J. L., Department of Plant Pathology, N.Y. State Agric. Expt. Station, Geneva, New York 14456. (315-787-2317)	(1)	USA
Noval Alonso, Cristina, Dept. de Proteccion Vegetal, Inst. Nacional de Investigaciones Agrarias, Apartado 8111, Madrid, Spain. (207–80–40, ext. 279)	(3)	SPN

Oberhofer, H., Sudtiroler Beratungsring fur Obst und Weinbau, A. Hoferstrasse 9, 390ll Lana, Sudtirol, Italy.	(3)	ITA
Okuse, I., Faculty of Agriculture, Laboratory of Hortic., Hirosaki University, Hirosaki, Aomori, Japan.	(3)	JAP
Olsson, Karen M., Swedish Univ. of Agric. Sciences, Dept. of Plant and Forest Protection, P. O. Box 7044, 750 07 Uppsala, Sweden. (018–10–20–00)	(3)	SWD
Opgenorth, D. C., Department of Plant Pathology, Univ. of California, Riverside, Calif. 92507. (714–787–4119)	(2)	USA
Otterbacher, A., University of Illinois, 105 Horticulture Field Laboratory, Urbana, Illinois 61801. (217–333–1520)	(2)	USA
Ottermann, A., Schering AG, Claudiusweg 13, 2077 Trittau, West Germany.	(2)	BRD
Paclt, J., Institute of Experimental Phytopathology and Entomology, Slovak Academy of Sciences, 900–28 Ivanka pri Dunaji, Czechoslovakia.	(3)	CZE
Paetzholdt, M., Pflanzenschutzamt, Hauptstrasse 108, 2084 Rellingen, West Germany.	(2)	BRD
Palazon, I., Departamento de Proteccion Vegetal, Centro de Investigaciones y Desarrollo Agrario del Ebro, Apartado 727, Zaragoza, Spain. (297207)	(3)	SPN
Panagopoulos, C. G., Benaki Phytopathological Institute, Kiffissia, Athens, Greece. (Ol-8013619)	(3)	GRC
Parnia, P., Scientific Director, Trustul Pomiculturii, Pitesti-Maracineni, Romania. (976–34.292)	(3)	ROM
Paulin, J. P., Station de Phytobacteriologie, I.N.R.A., Route de St. Clement, Beaucouze, 49000 Angers, France. (41-48.51.23)	(1)	FR
Pecknold, P. C., Department of Botany & Plant Pathology, Purdue University, West Lafayette, Indiana 47907. (317-749-6530)	(2)	USA
Persiel, F., Bundesforschungsanstalt für Gartenbauliche Pflanzenzuchtung, Bornkampsweg, 2070 Ahrensburg, West Germany. (04102–51122)	(1)	BRD
Petiot, J., Service de la Protection des Vegetaux, Cité Administrative, 59048 Lille Cedex, France. (20-52.12.21)	(1)	FR
Pilar, R., Departamento de Proteccion Vegetal, Centro de Invest. y Desarrollo Agrar. del Ebro, Apartado 727, Zaragoza, Spain (297207)	(3)	SPN

Porreye, W., Research Station of Gorsem, Brede Akker 3, 3800 Sint-Truiden, Belgium. (011-672019)	(1)	BLG
Preczewski, J. L., Product Development Dept., Stark Brothers Nurseries Co., Louisiana, Missouri 63353. (314–754–5009)	(2)	USA
Preiser, F., Research Laboratories, Merck and Company, Inc., Bldg. Rl23-l2, Rahway, New Jersey 07065. (201-574-6687)	(2)	USA
Prillwitz, H. G., Landespflanzenschutzamt, Essenheimerstr. 144, 6500 Mainz – Bretzenheim, West Germany.	(3)	BRD
<u>Psallidas</u> , P. G., Benaki Phytopathological Institute, Kiffissia, Athens, Greece. (Ol-8013619)	(3)	GRC
Quamme, H., Agriculture Canada, Research Station, Summerland, Brit. Columbia VOH 1ZO, Canada. (604–494–0401)	(2)	CND
Rackham, R. L., Benton County Extension Service, 2720 N.W. Polk Street, Corvallis, Oregon 97330. (503-776-7371)	(1)	USA
Reimann-Philipp, R., Bundesforschungsanstalt für Gartenbauliche Pflanzenzuchtung, Bornkampsweg, 2070 Ahrensburg, West Germany. (04102-51122)	(1)	BRD
Richter, J., Landesamt fur Pflanzenschutz, Reinsburgerstr. 107, 7000 Stuttgart-1, West Germany.	(2)	BRD
Ride, M., Station de Phytobacteriologie, I.N.R.A., Route de St. Clement, Beaucouze, 49000 Angers, France. (41–88.22.00)	(3)	FR
Ries, S. M., Department of Plant Path., Univ. of Illinois, N-427 Turner Hall, 1102 S. Goodwin, Urbana, Ill. 61801. (217-333-1523)	(1)	USA
Ristevski, B., Fruit Research Station, Faculty of Agric, 91000 Skopje, Yugoslavia (091–230–557)	(3)	YUG
Ritchie, D. F., Department of Plant Pathology, N. C. State University, Raleigh, North Carolina 27695. (919-737-2721)	(2)	USA
Roed, H., The Norwegian Plant Protection Institute, 1432 As-NLH, Norway.	(3)	NOR
Rom, R. C., Dept. of Hortic., Room 316, Plant Science Bldg. Univ. of Arkansas, Fayetteville, Ark. 72701. (501–575–2604)	(2)	USA
Roosje, G. S., Research Institute for Plant Protection, Binnenhaven 12, P. O. Box 9060, 6700 GW Wageningen, The Netherlands. (08370-19151, ext. 228)	(2)	NL

Rose, E., Hoechst AG, Landwirtsch. (Entwicklungsabteilung, Prufstelle Nord), Karl Wiechert Allee 3, 3000 Hannover 61, West Germany. (0511–5700.245)	(2)	BRD
Rosenberger, D. A., New York Agric. Exp. Station, Box 727 Highland, New York 12528. (914-255-8678)	(2)	USA
Ross, R. G., Agriculture Canada, Research Station, Kentville, Nova Scotia B4N 1J5, Canada. (902–678–2171)	(2)	CND
Rousselle, G. L., Agriculture Canada, Research Station, P. O. Box 457, St. Jean, Quebec J3B 6Z8, Canada. (514–346–4494)	(2)	CND
Rowson, G. R., Fruit Production Div., Showerings Vine Products and Whiteways Ltd., West Newton, Nr. Bridgewater, Somerset TA7 OBZ, England (412336)	(2)	UK
Rudolph, K., Institut fur Pflanzenpath. und Pflanzensch., Grisebachstr. 6, 3400 Gottingen, West Germany. (393721)	(2)	BRD
Russ, K., Bundesanstalt für Pflanzenschutz, Trunnerstrasse 5, 1021 Vienna, Austria. (0222–24.15.11)	(3)	OST
Ryugo, K., Department of Pomology, University of California, Davis, California 95616. (916–752–0929)	(2)	USA
Samson, Regine, Station de Phytobacteriologie, I.N.R.A., Route de St. Clement, Beaucouze, 49000 Angers, France. (41-87.69.97)	(2)	FR
Sanchezmonge, E., Departamento Genetica, Estac. Agronomos, Ciudad Universitaria, Madrid 3, Spain.	(3)	SPN
Sands, D. C., Dept. of Plant Path., Montana State Univ., Bozeman, Montana 59717. (406–994–4832)	(2)	USA
Sasser, M., Dept. of Plant Science, Univ. of Delaware, Newark, Delaware 19711. (302-738-2534)	(1)	USA
Schaper, U., Biologische Bundesanstalt, Institut fur Pflanzenbau im Obstbau, Postfach 73, 690l Dossenheim ub. Heidelberg, West Germany.	(2)	BRD
Scheer, H. A. T. van der, Research Station for Fruit Growing, Brugstraat 51, 4475 AN Wilhelminadorp, The Netherlands. (01100–16390)	(2)	NL
Schilli, E., Inst. fur Phytomedizin, Univ. Hohenheim, Otto-Sanderstrasse 5, 7000 Stuttgart-70, West Germany	(2)	BRD
Schmidle, A., Biologische Bundesanstalt, Institut für Pflanzenschutz im Obstbau, Schwabenheimerstrasse, Postfach 73, 690l Dossenheim/Heidelberg, West Germany. (06221-85238)	(3)	BRD

Schmidt, H., Pflanzenschutzamt des Landes Schleswig- Holstein, Westring 383, 2300 Kiel, West Germany.	(1)	BRD
Schroth, M. N., Department of Plant Pathology, Univ of California, Berkeley, Calif. 94720. (415–642–4147)	(1)	USA
Schulz, F. A., Inst. fur Phytopath., Christ.—Albrechts Univ., Olshausenstrasse 40–60, 2300 Kiel, West Germany. (0431–880–2996)	(1)	BRD
Schwabe, W. F. S., Fruit and Fruit Tech. Res. Inst., Private Bag 5013, 7600 Stellenbosch, South Africa (02231–2001)	(3)	SA
Seem, R. C., Department of Plant Pathology, N. Y. State Agric. Expt. Station, P. O. Box 462, Geneva, NY 14456. (315-787-2366)	(2)	USA
Seemuller, E., Biologische Bundesanstalt, Institut für Pflanzenschutz im Obstbau, Schwabenheimerstrasse, Postfach 73, 690l Dossenheim/Heidelberg, West Germany. (06221-85238)	(2)	BRD
Severin, V., Laboratory of Phytobacteriology, Research Inst. for Plant Protection, Blvd. Ion Ionescu de la Brad 8, Bucharest–Baneasa, Romania. (33.58.58–50)	(3)	ROM
Sharma, V. P., Dept. of Plant Pathology, Haryana Agric. University, Hissar 125004, India	(3)	IND
Simonsen, J., State Experimental Station, Laasbyvej 18, 8660 Skanderborg, Denmark. (06–520877)	(2)	DK
Slack, D., Dept. of Plant Pathology, Univ. of Arkansas, Fayetteville, Arkansas 72701. (501–575–2446)	(1)	USA
Smith, A. R. W., School of Biolog. Sciences and Environmental Health, Thames Polytechnic, Wellington Street, London SE18 6PF, England	(1)	UK
Sobiczewski, P., Research Institute of Pomology, ul. Pomologiczna 18, 96–100 Skierniewice, Poland. (Skierniewice 34–21)	(3)	POL
Soledad, S. V., Tagum, North Davao, Philippines 9401.	(3)	PHI
Spotts, R. A., Mid-Columbia Expt. Station, 3005 Expt. Station Drive, Hood River, Oregon 97031. (503-386-2030)	(2)	USA
Stankovic, D., Horticulture Dept., Faculty of Agriculture, Univ. of Belgrade, ul. Namanjina 6, 11080 Zemun (Belgrade), Yugoslavia.	(3)	YUG
Stark, C., Pflanzenschutzamt, Slevogtstr. 48, 2800 Bremen 1, West Germany.	(2)	BRD

Starr, M. P., Department of Bacteriology, University of California, Davis, Calif. 95616. (916–756–7000)	(1)	USA
Stead, D., Min. Agric. Fish. and Food, Harpenden Laboratory, Hatching Green, Harpenden, Herts AL5 2BD England (5241)	(1)	UK
Steiner, P., Department of Botany, University of Maryland, College Park, Maryland 20742. (301-454-3816)	(2)	USA
Stushnoff, C., Department of Horticultural Science, Univ. of Minnesota, St. Paul, Minnesota 55101. (612–373–1030)	(2)	USA
Sugar, D., Southern Oregon Expt. Station, 569 Hanley Rd., Medford, Oregon 97502. (503-772-5165)	(2)	USA
Suta, Victoria, Research Institute for Fruit Growing, 0300 Pitesti-Maracineni, Romania (976-34292)	(3)	ROM
Sutton, T. B., Department of Plant Pathology, N. C. State Univ., Raleigh, North Carolina 27650. (919–737–2752)	(1)	USA
Swanson, B. T., Dept. of Horticulture, 356 Alderman Hall, Universityof Minnesota, St. Paul, Minnesota 55108. (612–373–1011)	(1)	USA
Szkolnik, M., Department of Plant Pathology, N. Y. State Agr. Exp. Station, Geneva, New York 14456. (315–787–2375)	(1)	USA
Teissier, R., Service de la Protection des Vegetaux, 231 rue de la Convention, 75015 Paris, France. (532-21-29)	(3)	FR
Teylingen, M. van, Plant Protection Service, Geertjesweg 15, P. O. Box 9102, 6700 HC Wageningen, The Netherlands.	(2)	NL
Thibault, B., Station d'Arboriculture Fruitiere, I.N.R.A., Route de St. Clement, Beaucouze, 49000 Angers, France. (41-48.51.23)	(1)	FR
Thompson, J. M., USDA SE Fruit & Tree Nut Research Station, P. O. Box 87, Byron, Georgia 31008. (912-956-5656)	(1)	USA
Thomson, S. V., Department of Biology, Utah State Univ., Logan, Utah 84322. (801-750-3406)	(1)	USA
Travis, J. A., Department of Plant Pathology, Penn State University, Buckhart Lab., University Park, PA 16802.	(2)	USA
Tsiantos, J., Plant Protection Inst., Volos, Greece	(3)	GRC
Valyi, S., Department of Plant Protect. and Agrochemistry, Ministry of Agriculture and Food Admin., Kossuth Lajos ter 11, Budapest, Hungary.	(3)	HUN

Van Buskirk, P. D., Southern Oregon Expt. Station, 569 Hanley Rd., Medford, Oregon 97502 (503—772—5165)	(1)	USA
Vantomme, R., Laboratory for Microbiology, Ledeganckstraat 35, 9000 Gent, Belgium.	(1)	BLG
Veldeman, R., Ministry of Agriculture, Research Station for Phytopathology, Burg. van Gansberghelaan 96, 9220 Merelbeke, Belgium. (091–522083)	(2)	BLG
Vereecke, M., Belgian Plant Protection Service, Rue de Stassart 36, 1050 Brussels, Belgium	(2)	BLG
Vogelsanger, D., Pflanzenschutzamt, Hermannswerder 20A, 15 Potsdam, East Germany.	(3)	DOR
Vondracek, J., Fruit Research Station, Techobuzize, 411 42 Ploskovice (okr. Litomerice), Czechoslovakia. (Ploskovice 9387)	(3)	CZE
Voronkova, L., Dept. of Bacteriology, Central Laboratory for Plant Quarantine, 1/11 Orlikov per., 107139 Moscow, B-139, Russia.	(3)	RUS
Vukovits, G., Bundesanstalt für Pflanzenschutz, Vienna, Austria.	(3)	OST
Wade, E. K., Department of Plant Pathology, University of Wisconsin, Madison, Wisconsin 53706. (608–262–1426)	(2)	USA
Waldner, W., Sudtiroler Beratungsring fur Obst und Weinbau, Andreas Hoferstrasse, 39011 Lana, Italy	(3)	ITA
Walsh, P. F., Dept. of Agriculture, Agriculture House, Kildare St., Dublin 2, Ireland. (789011, ext. 3289)	(3)	IRL
Way, R. D., Department of Pomology & Viticulture, N. Y. State Agric. Expt. Station, Geneva, New York 14456. (315–787–2235)	(1)	USA
Westwood, M. N., Department of Horticulture, Oregon State University, Corvallis, Oregon 97331. (503-754-3695)	(2)	USA
Wiggel, D., Ministry of Agriculture, Olantigh Road, Wyl, Nr. Ashford, Kent, England.	(2)	UK
Willett, M., Extension Service, Oregon State University, 1301 Maple Grove Drive, Medford, Oregon 97501. (503-776-7371)	(2)	USA
Wimalajeewa, S., Plant Research Institute, Burnley Gardens Swan Street, Burnley, Victoria 3121, Australia. (8101511)	(3)	AUS

Wodzinski, R. S., Biology Department, Ithaca College, Ithaca, New York 14850. (607–274–3979)	(1)	USA
Yoder, K. S., Fruit Research Laboratory, Va. Polytech. Inst., 2500 Valley Ave., Winchester, Virginia 22601. (703–667–8330)	(1)	USA
Yorston, Y. M., Brit. Columb. Ministry of Agric., Research Station, Summerland, British Columbia VOH 1ZO, Canada. (604–494–0401)	(2)	CND
Young, J. M., Plant Diseases Division, MARC, Private Bag, Auckland, New Zealand. (893660)	(2)	NZ
Zehr, E. I., Department of Plant Pathology & Physiology, Clemson University, Clemson, South Carolina 29631. (803–656–3450)	(2)	USA
Zeller, W., Biologische Bundesanstalt für Land und Forstwirtschaft, Institut für Pflanzenschutz in Ackerbau und Grunland, Schlosskoppelweg 8, 2305 Heikendorf-Kitzeberg (Kiel), West Germany. (0431-23495)	(1)	BRD
Zoller, B. G., The Pear Doctor, Inc., P. O. Box 952, Yuba City, California 95991. (916-674-1255)	(2)	USA
<pre>Zwet, T. van der, U. S. Department of Agriculture, Appalachian Fruit Research Station, Rt. 2, Box 45, Kearneysville, West Virginia 25430. (304-725-3451, ext. 29)</pre>	(1)	USA

Working Group Membership by Country $\frac{1}{2}$

Argentina	Bergna, D. A. Dobra, A.	*Meyer, F. C.
Australia	<pre>#Cartwright, D. N. Wimalajeewa, S.</pre>	
Austria	Russ, K. Vukovits, G.	
Belgium	Deckers, T. De Ley, J. Geenen, J. Laere, O. van Laroche, M. Luchene, K. van	Maroquin, C. *Porreye, W. Vantomme, R. Veldeman, R. Vereecke, M.
Brazil	Bredemeier, D. Feliciano, A. J.	
Canada	Agr. Can. Library Biggs, A. R. *Bonn, W. G. Cline, R. A. Coulombe, L. J. Crowe, A. D. Davidson, J. G. N. *Evans, I. R. Gibbins, L. N. Horricks, J.	Howard, R. J. Hunter, C. L. Kappel, F. Lane, D. Lethal, J. *McPhee, R. Muir, J. Quamme, H. *Ross, R. G. Rousselle, G. L. Yorston, Y. M.
China (P.R.)	*Cao, R.	
Czechoslovakia	*Kudela, V. Paclt, J.	Vondracek, J.
Denmark	Andersen, H. Christensen, F. G. *Dinesen, G. Jorgensen, H. A.	Kristensen, H. R. Mosegaard, J. Simonsen, J.
East Germany	*Kleinhempel, H. Muller, H. J.	Vogelsanger, D.
Egypt	*Abo-El-Dahab, M. K. Mickail, K. Y.	

 $[\]frac{1}{N}$ Names with asterisk (*) are contact persons.

England	Alston, F. H. Bennett, R. A. Billing, E. Byrde, R. J. W. Cooper, R. M. Fox, R. T. V. *Garrett, C. M. E.	Gwynne, G. Hignett, R. C. Lelliott, R. A. Rowson, G. R. Smith, A. R. W. Stead, D. Wiggel, D.
France	Balavoine, P. Callu, D. Large, M Lecomte, P. Mathys, G. *Paulin, J. P.	Petiot, J. Ride, M. Samson, R. Teissier, R. Thibault, B.
Greece	Panagopoulos, C. G. *Psallidas, P. G.	Tsiantos, J.
Hungary	*Klement, Z. Valyi, S.	
India	Gupta, V. K.	Sharma, V. P.
Ireland	*Walsh, P.	
Italy	*Bazzi, C. Calzolari, A. Ercolani, G. L. Fideghelli, C.	Garibaldi, A. Mazzucchi, U. Oberhofer, H. Waldner, W.
Japan	Goto, M. Kato, T.	*Okuse, I.
Могоссо	Benjama, A.	⇔Chouibani, M.
Mexico	*Fucikovsky, L.	Mendoza, H., A.
<u>Netherlands</u>	Bouma, S. CHRONICA HORTIC. Heybroek, H. M. Kooistra, T. Langeslag, J. J. J. *Maas Geesteranus, H. P.	Mijneke, C. A. R. Miller, H. J. PUDOC Roosje, G. S. Scheer, H. A. T. van der Teylingen, M. van
New Zealand	Dye, D. W. *Hale, C. N.	Young, J. M.
Norway	Dale, T. *Roed, H.	
Philippines	Soledad, S. V.	

Burkowicz, A. Poland *Sobiczewski, P. Conceicao, J. *Martins, J. M. S. Portugal *Suta, V. Romania Parnia, P. Severin, V. *Voronkova, L. Russia South Africa Button, J. *Matthee, F. N. Erskine, J. M. Schwabe, W. F. S. Spain Lopez, Gonzalez, M. Palazon, I. Mansergas, A. J. F. Pilar, R. *Noval Alonso, C. Sanchezmonge, E. *Graberg, M. Olsson, K. M. Sweden Kroeker, G. Switzerland Bolay, A. *Grimm, R. Cazelles, O. Joseph, E. Egli, T. Baykal, N. Turkey West Germany Baumm, L. H. Muller, K. Brulez, W. Ottermann, A. Cornils, H. Paetzholdt, M. Persiel, F. Duben, J. Franz, W. Prillwitz, H. G. Graf, H. Reimann-Philipp, R. Hoppe, H. Richter, J. Isenbeck, M. Rose, E. Knosel, D. Rudolph, K. Schaper, U. Kraus, P. Kuhne, H. Schilli, E. Lehmann-Danzinger, H. Schmidle, A. Lux-Wellenhof, E. Schmidt, H. Mappes, D. Schulz, F. A. Massfeller, D. *Seemuller, E. Meyer, J. Stark, C. Michel, H. G. *Zeller, W.

Arsenijevic, M.

Ristevski, B.

*Stankovic, D.

Yugoslavia

USA

Abdel-Rahman, M. Aldwinckle, H. S. Lombard, P. B. Ark, P. A. Luepschen, N. S. *Barrat, J. G. McSwan, I. C. Bates, J. J. *Beer, S. V. *Miller, R. W. Morehead, G. W. Bell, R. L. Morton, H. V. Berry, D. W. Norelli, J. L. Beutel, J. A. Opgenorth, D. C. Biehn, W. Burr, T. J. Otterbacher, A. Pecknold, P. C. Bushong, J. W. Preczewski, J. L. Cameron, H. R. *Preiser, F. Carlson, R. F. Rackham, R. L. Carroll, V. J. *Ries, S. M. Chandler, D. *Ritchie, D. F. Civerolo, E. L. Rom, R. C. Clayton, C. N. Rosenberger, D. A. *Covey, R. P. Ryugo, K. Crassweller, R. Sands, D. C. Cummins, J. N. Sasser, M. Davidson, S. Schroth, M. N. *Douglas, S. M. Seem, R. C. *Drake, C. R. *Slack, D. Egolf, D. R. Spotts, B. P. *Ellis, M. A. Starr, M. P. French, J. R. *Steiner, P. Gantotti, B. V. Stushnoff, C. Gates, D. *Sugar, D. *Goodman, R. N. Sutton, T. B. Swanson, B. T. Harnish, W. Heimann, M. F. Szkolnik, M. *Hickey, K. D. *Thompson, J. M. Hildebrand, E. M. *Thomson, S. V. Travis, J. A. *Janick, J. Van Buskirk, P. D. Johnson, D. E. Jones, A. L. *Wade, E. K. Kado, C. I. Way, R. D. *Klos, E. J. Westwood, M. N. Koenigshof, R. Willett, M. Kuc, J. Wodzinski, R. S. Yoder, K. S. Kyle, N. E. Lacy, G. H. Zehr, E. I. Lamb, R. C. *Zoller, B. G. Landis, W. R. Zwet, T. van der

SUMMARY

Contact Persons for Fire Blight Newsletter

United	States	Oth	er Countries
Arkansas	Slack, D.	Argentina	Meyer, F. C.
California	Zoller, B. G.	Australia	Cartwright, D. N.
Connecticut	Douglas, S. M.	Belgium	Porreye, W.
Georgia	Thompson, J. M.	China (P.R.)	Cao, R.
Illinois	Ries, S. M.	Czechoslovakia	Kudela, V.
Indiana	Janick, J.	Denmark	Dinesen, A.
Maryland	Steiner, P.	Egypt	Abo-El-Dahab, M.K.
Michigan	Klos, E. J.	England	Garrett, C. M. E.
Missouri	Goodman, R. N.	France	Paulin, J. P.
New Jersey	Preiser, F.	Germany (East)	Kleinhempel, H.
New York	Beer, S. V.	Germany (West)	Seemuller, E.
North Carolina	Ritchie, D. F.		Zeller, W.
Ohio	Ellis, M. A.	Greece	Psallidas, P. G.
Oregon	Sugar, D.	Hungary	Klement, Z.
Pennsylvania	Hickey, K. D.	Ireland	Walsh, P.
South Carolina	Miller, R. W.	Italy	Bazzi, C.
Utah	Thomson, S. V.	Ja pan	Okuse, I.
Virginia	Drake, C. R.	Mexico	Fucikovsky, L.
Washington	Covey, R. P.	Morocco	Chouibani, M.
West Virginia	Barrat, J. G.	Netherlands	Maas Geesteranus, H. P.
Wisconsin	Wade, E. K.	New Zealand	Hale, C. N.
		Norway	Roed, H.
		Poland	Sobiczewski, P.
		Portugal	Martins, J. M. S.
		Romania	Suta, V.
Canad	da	Russia	Voronkova, L.
		South Africa	Matthee, F. N.
Alberta	Evans, I. R.	Spain	Noval Alonso, C.
British Columbia	McPhee, R.	Sweden	Graberg, M.
Nova Scota	Ross, R. G.	Switzerland	Grimm, R.
Ontaria	Bonn, W. G.	Yugoslavia	Stankovic, D.

SUMMARY

Persons Interested in Fire Blight

	I	nterest	Category			Number of Contact
Country	1	2	3	4	Total	Persons
* USA - United States	37	48		4	89	21
* CND - Canada	3	18			21	4
<pre># BRD - West Germany</pre>	12	19	3		34	2
* UK - England	9	5			14	1
* NL - Netherlands	5	7			12	1
* FR - France	5	3	3		11	1
* BLG - Belgium	7	4			11	1
* DK - Denmark	1	6			7	1
* DDR - East Germany			3		3 3 2	1
NZ - New Zealand	1	2			3	1
* POL - Poland	1		1		2	1
* EGY - Egypt	2				.2	1
* MEX - Mexico		1			1	1
ITA - Italy			8		8	1
SPN - Spain			6		6	1
SWT - Switzerland			5		5	1
SA - South Africa			4		4	1
ARG - Argentina			3		3	1
CZE - Czechoslovakia			3 3 3 3 3 3 2 2 2		3 3 3 3 3 2 2 2 2 2 1	1
JAP - Japan			3		3	1
ROM - Romania			3		3	1
SWD - Sweden			3		3	1
YUG - Yugoslavia			3		3	1
GRC - Greece			3		3	1
AUS - Australia			2		2	1
HUN - Hungary			2		2	1
POR - Portugal			2		2	1
NOR - Norway			2		2	1
MOR - Morocco			2		2	1
IRL - Ireland			1			1
CHI - China			1		1	1
RUS - Russia			1		1	1
OST - Austria			2 2 2		2 2 2 1	
BRA - Brazil			2		2	
IND - India					2	
PHI - Philippines			1			
TUR - Turkey			1		1	
TOTAL	83	113	75	4	275	56

^{*}Countries with fire blight.

		,	
:			

Fire Blight Mailing List Questionnaire

*

*

*

*

*

* * *

*

The list of names in this Newsletter is an annual attempt to establish a complete and updated mailing list of all persons interested in fire blight. Please make corrections and additions where necessary and send me any new names not listed. A new list will be prepared for the next newsletter.

	My name, address and telephone are correct (if not, show change below)
	My interest in fire blight is correct (if not, please indicate below)
	My name should be dropped from this list
	My/other name should be added to this list
NAME	
ADDRESS	
	ZIP
TELEPHONE	
Interest in fire bligh	ot research: 1 2 3 4
Interest in fire bligh	nt newsletter: YES NO Please circle
I will serve as contact for newsletter ques	t person

Please return to your contact person or directly to:

T. van der Zwet Appalachian Fruit Research Station Route 2, Box 45 Kearneysville, West Virginia 25430



